



京都大学大学院医学研究科 てんかん・運動異常生理学講座

Department of Epilepsy, Movement Disorders and Physiology
Kyoto University Graduate School of Medicine

年次報告書
Annual Report 2016

2016年8月

<表紙の言葉>

表紙のデザインは、波形様の曲線と、てんかんキャンペーンの紫と多彩な色からなります。脳波の波形と周波数を想像させます。脳波のサーフィンがもっと上手になることを目指して。

<Front cover>

Design of the cover page is the slow waveforms with different colors including purple, which may remind you of brain waveforms and different frequencies. Hoping all of you to enjoy EEG wave surfing.


**京都大学大学院医学研究科
てんかん・運動異常生理学講座**

Department of Epilepsy, Movement Disorders and Physiology
Kyoto University Graduate School of Medicine

**年次報告書
Annual Report 2016**

2016年8月

Preface



Since two years have past after we have issued the first and second annual reports of the Department of Epilepsy, Movement Disorders and Physiology, Kyoto University Graduate School of Medicine, it is again my great honor and pleasure that we could issue the third annual report. We appreciate very much all of you who kindly gave the very important comments and suggestion, upon the previous annual reports from other places and fields.

First of all, since the establishment of this department in August 2013, we have been supported, as the endowed department, by GlaxoSmithKline K.K., Niho Kohden Co., Otsuka Pharmaceutical Co., and UCB Japan Co. Ltd. Within the Kyoto University, we greatly appreciate the warmest and continuous support by the Department of Neurology (chaired by Prof. Ryosuke Takahashi), and also by the Departments of Neurosurgery (Prof. Susumu Miyamoto), Psychiatry (Prof. Toshiya Murai), Pediatrics (Prof. Toshio Heike), and Diagnostic Imaging & Nuclear Medicine (Prof. Kaori Togashi).

Based on or in addition to the daily clinical practice for patients with epilepsy and movement disorders in the Kyoto University Hospital, the aim of this Department was originally planned as follows.

- 1) Clinical Research: To solve pathophysiology of epilepsy and movement disorders, and to develop new treatments that could lead to the advanced medical care. We also aim at standardizing it in clinical practice.
- 2) Translatable and Translational Research: To promote clinical practice and research of clinical neurophysiology on epilepsy and movement disorders and its clinical application, because basic- and clinical epileptology and movement disorders are very closely related to neurophysiological knowledge and methods.
- 3) Education: To provide the integrated teaching opportunity for training of physician-scientists, clinical specialists and leaders of related fields internationally.
- 4) Industry-academia collaboration under the appropriate COI management

Upon the completion of the third year, we again greatly appreciate all of the Departments of Kyoto University Graduate School of Medicine and all of the Clinical Divisions of the Kyoto University Hospital for their warmest and sincere support to our activity. Without this supports, we could not complete any of them by all means that were reported in this annual report. It was also strongly supported by so many collaborators, researchers and friends in Japan and also internationally.

In this annual report, we are also very pleased to report the outcome and the current state of industry-academia collaboration. We hope that it may represent a positive face of endowment department in general and that it may help industry partners understand the significance of the endowment department in the future plan.

We again tried to summarize what we could do and not, which help us analyze and understand the situation, and then help us improve and modify the current condition, and also start the new concerns. We would greatly appreciate your any feedback to us that is very helpful for our future contribution to the patients close to or distant from us.

At the end of my preface, I thank all of our friends very much whoever kindly worked hard to make and edit this annual report.

August 2016

With my best wishes,

A handwritten signature in black ink, appearing to read "Akio Ikeda".

Akio IKEDA, MD, PhD
Chairman and Professor
Department of Epilepsy, Movement Disorders and Physiology
Kyoto University Graduate School of Medicine

Congratulatory remarks



On behalf of the Vice President of International League Against Epilepsy (ILAE), I want to congratulate Professor Akio Ikeda, for his 3rd year Anniversary Remark of Department of Epilepsy, Movement Disorders and Physiology, Kyoto University Graduate School of Medicine. First of all, I also want to congratulate Professor Akio Ikeda, for his great efforts to realize and perform such outstanding programs, for clinical and basic researches, education, administration, as well as clinical treatments for patients with neurological disorders.

Professor Akio Ikeda is outstanding neurologist and epileptologist not only in Japan and Asian friend countries but also in all over the world-wide countries.

In ILAE, publication of the research articles for Official Journal (Epilepsia) is one of the most important activities. Professor Akio Ikeda was nominated as an Associate Editor of the Epilepsia since 2013, and he not only organized and controlled many reviewers with his amazing abilities but also he reviewed many articles whether to accept or reject for Epilepsia with his sharp decision. I am very happy to see his contribution because I heard many reputations of his good work from ILAE Executive members. I can always remember his earnest but strict attitude for clinical and basic researches.

Recently, World Health Organization (WHO) approved 'Global burden of Epilepsy and the need for coordinate action at the country level to address its health, social and public knowledge implications', at WHO General assembly in Geneva, in February 2015. This decision is now a world-wide movement for epilepsy to introduce and implement national health care plans of action for epilepsy management, aiming to overcome inequalities and inequities in health, social and other related services including Japan. I am sure that Professor Akio Ikeda can further develop this important resolution for patients with epilepsy in coming next step in his department.

With my best wishes,

A handwritten signature in black ink, appearing to read "Tatsuya Tanaka".

Tatsuya Tanaka, MD., DMSc.
Vice President, ILAE 2009-2017
Past President, Japan Epilepsy Society
Professor Emeritus, Asahikawa Medical College
Honorary Director, Yamabiko Meidcal Welfare Center

Index

I. Introduction	47
II. Outline	49
III. Activity report	51
Research activities	51
Clinical activities	61
Educational activities	64
Research grants obtained from extramural sources & awards	67
IV. Society-academia collaboration	69
V. Publications and Congress Presentations	27
Please see the contents in the Japanese section of the annual report (pages 27-40).	
Publications	27
Original articles	27
Edited books & Journal	29
Book chapters	29
Review papers	31
Presentations	33
Congress presentations	33
Invited lectures and symposium etc.	33
Oral and poster presentations	34
Other presentations	37
Educational lectures	39
Intramural lectures	40
VI. Attached materials	71
Collaborative research activity in the Grant-in-Aid for Scientific Research on Innovative Areas: Non-linear Neuro-oscillology - Towards Integrative Understanding of Human Nature	71
9th electroencephalography (EEG) and electromyography (EMG) seminar	72
1st advance course of electroencephalography (EEG) seminar	73
104th Kinki Regional Division of Japanese Society of Neurology	73
Epilepsy Lectures in Autumn 2015, Kyoto	74
5th Masakazu Seino Memorial Lecture	74
Visiting physicians	75

I Introduction

To begin with

The details of the annual report in each aspect are shown in the following pages as done in the second annual report last time, and thus as the essence of general message directly related to our Department, I am briefly listing them which occurred or appeared in the last year. I would greatly appreciate your comment and feedback.

- 1) Evolutional changes in the social state for epilepsy
 - 2) "Oscillology": the new research field including epilepsy in Japan
 - 3) EEG/ Epilepsy fellowship in Kyoto University Hospital
 - 4) Teaching session
 - 5) Epilepsy should be approached based on the broad scope of general neurology and clinical neuroscience.
-
- 1) Epilepsy used to be recognized as a neuropsychiatric disease in Japan historically. Once recently the genetic study, neurophysiological analysis and pathological approach have advanced better understanding of epilepsy, it is now recognized one of the most representative disorders of clinical neuroscience for both neurology and neuropsychiatry. As shown by the fact that World Health Assembly closes, passing resolutions on epilepsy. <http://www.who.int/mediacentre/news/releases/2015/wha-26-may-2015/en/>. It has aided in improving clinical care in epilepsy in Japan as follows. Insurance reimbursement in epilepsy has greatly improved since last April 2016 by the National Healthcare System for prolonged-video EEG monitoring, remote reading system of digital EEG and EEG reading service.
 - 2) With regard to clinical and translatability research, we have been appointed as the member of Grant-in-Aid for Scientific Research on Innovative Areas (Research in a proposed research area): Non-linear Neuro-Oscillology - Towards Integrative Understanding of Human Nature. We will aim at exploring novel multi-dimensional oscillatory phenomena, working mainly on primate/rodent models, direct recordings of human brain activity, and advanced measurements of human brain systems. (<http://www.nips.ac.jp/oscillology>)
 - 3) We have provided the special EEG/ Epilepsy fellowship as the joint activity with the Department of Neurology (Prof. Takahashi), and 1-3 fellows have been always employed so far. As for clinical care in the third year, more than that of the first and second year, we tried to contribute to the activity of the Kyoto University Hospital and Graduate School of Medicine, as much as possible, by conducting our initial purposes.
 - 4) We also tried to contribute to the region of Kyoto and Kansai directly by patient care, and also tried to provide the teaching seminars in the field of clinical EEG, clinical epilepsy, and clinical neurophysiology by means of many medical societies, i.e., Japanese Society of Neurology, Japanese Society of Clinical Neurophysiology, and Japan Epilepsy Society.
 - 5) Epilepsy should be approached based on the broad scope of general neurology and clinical neuroscience. So-called autoimmune epilepsy is one of the most important topics in clinical care and research in epilepsy, and neuroimmunological approaches are very important from general neurology's point of view. Glia has also been regarded very important for both neurodegenerative disorders and epilepsy. Epilepsy is one of the commonest neurological disorders, and thus approaches to epilepsy in patient's care, research and teaching should be done based on the broad scope of general neurology and clinical neuroscience.

I. Introduction

In the last, for young people, we very much welcome all of you who are interested in this field from any places, and we hope to work and learn together.

With my best wishes,

Akio IKEDA, MD, PhD



II Outline

Funding prospects

Establishment	June 1st, 2013
Arrival of staffs	August 1st, 2013

Name of the Endowed Department

Department of Epilepsy, Movement of Disorders and Physiology
Kyoto University Graduate School of Medicine

Founding vision

- 1) As an academic department in the university hospital, we promote researches and clinical applications of clinical neurophysiology, which is essential for elucidating the pathophysiology and developing the treatment of clinical epileptology.
- 2) It is our mission to elucidate the pathophysiology and develop the treatment of epilepsy and movement disorders so as to develop the highly advanced medicine for its application to clinical practice. We also make best efforts to offer opportunities of trainings and educations to young physicians both from Japan and abroad to foster the specialists and physician-scientists in the field of epilepsy.

Research vision

In close collaboration with the Department of Neurology, we aim at achieving the following projects in a comprehensive, efficient and multidisciplinary manner.

1. Development of medical devices for wide-band EEG recording&analysis, and its application to elucidation of epileptogenicity
2. Promotion of epilepsy surgery and research on higher brain functions&its plasticity under epileptic conditions
3. Combined imaging and neurphysiological researches on the pathophysiology of the epileptic focus
4. Research on the pathophysiology and treatment of movement disorders
5. iPS (induced pluripotent stem) cell research and animal model research on epileptogenesis
6. Establishment of the training programs for the advanced specialists in the related fields
7. Promote collaborative researches with basic and mathematical scientists to understand neural oscillations underlying both physiological brain functions and pathology

Companies of endowment (in alphabetical order)

GlaxoSmithKline K.K.
NIHON KOHDEN CORPORATION
Otsuka Pharmaceutical Co., Ltd.
UCB Japan Co., Ltd.

Contact address

URL: <http://epilepsy.med.kyoto-u.ac.jp>
E-mail: epilepsy@kuhp.kyoto-u.ac.jp (Secretary)
Address: Shogoin, Sakyo-ku, Kyoto, 606-8507, JAPAN
TEL: (+81)-75-751-3662
FAX: (+81)-75-751-3663

II. Outline

Members of this Department and Affiliated Persons

<Members of Department of Epilepsy, Movement Disorders and Physiology>

Professor: Akio Ikeda, M.D., Ph.D. Associate Professor: Riki Matsumoto, M.D., Ph.D. (- Jul. 2016)
Secretary: Miki Watanabe Medical assistant technician: Makiko Matsushita (Apr. 2016 -)
Visiting Scientist: Kiyohide Usami, M.D., Ph.D. (Dept. of Neurology, Rakukakai Otowa Hospital, Nov. 2014 - Sep. 2015)
Tomoyuki Fumuro, Ph.D. (Dept. of Medical Technology and Sciences, School of Health Sciences at
Fukuoka, International University of Health and Welfare, Mar. 2015 -)
Research support personnel: Takeshi Inoue M.D. (Dept. of Pediatric Neurology, Osaka city general hospital, May 2016 -)

<Affiliated members from Department of Neurology>

Associate Professor: Riki Matsumoto, M.D., Ph.D.
(Aug. 2016 -)
Assistant Professor: Akihiro Shimotake, M.D., Ph.D.
Katsuya Kobayashi , M.D., Ph.D.

Graduate Students (Doctoral course):

M. Ota, Kei. Sato, H. Takeyama, M. Daifu-Kobayashi,
M. Kinboshi (from Wakayama Medical University), T. Murai,
M. Nakatani (from Juntendo University School of Medicine),
J. Togawa, K. Yoshinaga, S. Neshige (from Hiroshima
University School of Medicine), Kazuaki. Sato, K. Tanioka,
M. Togo, M. Sakamoto

Graduate Student (Master's course): Shamima Sultana

EEG/Epilepsy fellowship:

T. Inoue (Apr. 2013 – Mar. 2016)
H. Yoshimura (Jul. 2015 – Sep. 2015)
T. Tsukada (Oct. 2015 – Mar. 2016)
D. Fujii (Feb. 2016 -)
M. Honda (Apr. 2016 -)

Research student and undergraduate student:

B. Bayasgalan (Apr. 2016 -)
R. Kurauchi
(Department of Human Health Sciences, Apr. 2016 -)

<Affiliated members>

Department of Neurosurgery

Senior Lecturer: T. Kunieda, M.D., Ph.D.
(- Apr. 2016, Ehime University)
K. Yoshida, M.D., Ph.D.
Assistant Professor: T. Kikuchi, M.D., Ph.D.

Human Brain Research Center Research and Educational Unit of Leaders for Integrated Medical System

Program-specific Associate Professor:
M. Matsuhashi, M.D., Ph.D.

Department of Clinical Laboratory Medicine

Assistant Professor: T. Hitomi, M.D., Ph.D.

Department of Respiratory Care and Sleep Control Medicine

Assistant Professor: M. Inouchi, M.D., Ph.D.





Activity report

Research activities

Our main goal is to solve so-called "clinical questions", which have been raised in the daily clinical activity, and have remained unsolved yet.

Based on the concept of systems neuroscience, by means of established and newly developed various methods, many clinical and basic researches are conducted as follows.

KEY WORDS

General key words: epilepsy, epilepsy surgery, higher cortical function (motor control, praxis, language, semantic cognition, vision, will), Bereitschaftspotential (BP), cortico-cortical network, movement disorders, sleep disorders, autoimmune epilepsy

Unique key words: ictal DC shift, high frequency oscillation (HFO), cortico-cortical evoked potential (CCEP), cortical tremor, ictal apraxia, ictal paresis, wide-band EEG

1) Pathophysiology of medically intractable epilepsy and its treatment

a) Unraveling pathophysiology for intractable partial epilepsy

Epilepsy surgery has been established as an option for treatment of intractable partial epilepsy. The epileptogenic lesions, such as hippocampal sclerosis, cavernous angioma, and brain tumor, are the most common candidates for the one-stage surgery. It is still a challenge to localize the epileptic focus in 'MRI-negative' patients, who show no (apparent) lesion by clinical MRI studies. We have extensively made multidisciplinary approaches with video-EEG monitoring, MEG and FDG-PET to 'visualize' the epileptic focus in these MRI-negative patients for possible treatment with epilepsy surgery. Especially in the MRI-negative cases, we occasionally need invasive evaluation with intracranial electrodes to precisely delineate the epileptic focus and map the eloquent cortices at and around the epileptic focus. In addition to the conventional frequency band (Berger rhythm: 0.3-70 Hz), advancement in medical engineering has enabled us to record electrocorticogram with a broader frequency band (wide-band EEG), ranging from direct current shifts (DC shifts) to high frequency oscillations (HFOs). In our recent report, several features of ictal DC shifts and HFOs were extracted in 16 patients who underwent chronic intracranial recording as follows; 1) high occurrence rates of DC shifts during ictal period, 2) DC shifts preceded HFOs with statistically significance in most patients, 3) DC shifts and HFOs were observed in more restricted area than conventional ictal pattern, 4) higher occurrence rate in focal cortical dysplasia than hippocampal sclerosis (Kanazawa et al., 2014). We have been applying this wide-band EEG technology (ictal/interictal DC shift and HFOs) to further localize the core epileptogenic zone for epilepsy surgery. Furthermore, we started collaborative studies with Osaka University of Pharmaceutical Sciences to elucidate generation mechanisms of DC shifts and HFOs by means of wide-band EEG analysis and multi-unit recording in epilepsy model animals.

It has been known that epileptic discharges can spread to the other areas through intercortical networks. Thus, in addition to epileptic foci, it would be important to understand an epileptic network formulated by epileptic foci, propagating areas and connecting white matters. MRI tractography is a method evaluating anatomical networks. By combination with MR tractography and FDG-PET, we delineated decreased integrity of seizure propagation tracts in MTLE patients (Imamura et al., 2016). We recruited 18 patients with MTLE and 18 healthy subjects. In the patients, the remote functional deficit zone was defined using FDG-PET as an extratemporal region showing glucose hypometabolism. Using diffusion MRI tractography, we defined a seizure propagation tract (PT) that connects the focus with a remote functional deficit zone. We also used the corticospinal tract (CST) and inferior longitudinal fasciculus (ILF) as control tracts. Fractional anisotropy (FA), mean diffusivity (MD), and volume of the tracts were compared among PT, CST, and ILF. Tractographic analysis identified the uncinate fasciculus, arcuate fasciculus, and fornix as PTs. A decrease in FA was found in MTLE patients compared with healthy subjects in all tracts, but




Figure 1. (Center) We delineated a seizure propagation tract (PT) that connects the focus with a remote functional deficit zone. PT goes through mainly uncinate fasciculus (UF). We also used the corticospinal tract (CST) and inferior longitudinal fasciculus (ILF) as control tracts.

(Left) Whole brain analysis using TBSS delineated a widespread decreased FA in the MTLE patient group, which involved the PT, ILF and CST.

(Right) A decrease in FA was found in MTLE patients compared with healthy subjects in all tracts, but PTs showed a more significant decrease in FA than did the two control tracts (modified from Imamura et al., 2016).

PTs showed a more significant decrease in FA than two control tracts (Fig. 1). Although white-matter damage was observed in all candidate tracts examined, the integrity of white matter was most significantly decreased in PTs in MTLE. The present findings demonstrate that epileptic activity impairs not only the focus, but also the whole network (the focus - the propagation tract - the remote functional deficit zone).

Cortico-cortical evoked potential, CCEP, is an *in vivo* electrical tract tracing method for presurgical evaluation, developed in Cleveland Clinic and Kyoto University (Matsumoto et al., Brain 2004). By means of subdural electrodes, electrical pulses (0.3-ms duration, frequency of 1 Hz, alternating polarity, 1–10 mA) are directly delivered to a part of the cortices, and CCEPs are obtained from adjacent and remote cortical regions by averaging the electrocorticogram time-locked to the stimulus onset. It promises to refine our understanding of surgical candidacy, first through a more precise and tailored evaluation of the seizure network in each individual patient, and second through greater understanding of the functional systems of the brain involved. Both are important for improving our ability to identify patients at high risk for poor surgical outcomes across multiple outcome measures.

b) Unraveling pathophysiology and treatment for various epilepsy syndrome

Even in the 21st century, electroencephalography (EEG) remains essential in the diagnosis of "epileptogenicity". Simultaneous recording of EEG and fMRI (EEG-fMRI) is a new technique that takes advantages of both modalities and complements each other to delineate both the cortical and subcortical structures related with epileptic activities. We for the first time introduced this technique to Japan to investigate the epileptic network and underlying pathophysiology in various types of epileptic syndromes such as praxis-induced epilepsy and hypothalamic hamartoma (HH). Recently, we demonstrated epileptic networks of HH by means of EEG-fMRI (Usami et al., 2016). Eight patients with HH were recruited, and underwent EEG-fMRI. A general linear model was utilized for analyzing fMRI, using time-shift models designed by timing at and around spikes (from -8 sec to + 4 sec) convolved with hemodynamic response function. Any BOLD responses were observed in all patients, and six out of eight patients

presented BOLD responses at/around the hypothalamus ipsilateral to attachment of the hamartoma. In group analysis, activations included the ipsilateral hypothalamus, brainstem tegmentum, bilateral precunei, lateral parietal lobes and the contralateral dominant bilateral cerebellum, and deactivations included the ipsilateral cuneus, bilateral caudate nuclei, thalamus, hippocampi, and a part of default mode network (DMN) (Fig. 2). Based on the results, hypothalamus, brainstem, cerebellum and cerebral cortices would be a network generating epileptic activities, and DMN and hippocampi might be relevant to loss of awareness during seizures and epileptic encephalopathy in HH patients.

Recently the role of autoimmunity has been highlighted for a subset of encephalitis and epileptic seizures. Nowadays, high-resolution MRI can depict subtle structural changes in the medial temporal lobe, such as hippocampus and amygdala, which are often the main inflammatory foci in autoimmune limbic encephalitis. By




Figure 2. Group analysis of EEG-fMRI. Group analysis of seven patients revealed the common epileptic network associated with epilepsy with HH. Before group analysis, MRI images were flipped in the left-right dimension so that the right hemisphere shows the side of attachment (unilateral attachment) or that of predominant hypothalamic activation (bilateral attachment). The voxels with significant activation/deactivation by cluster-level statistics ($p < 0.007$ after Bonferroni correction) are overlaid on the group mean anatomical image in the MNI standard space. Activations included the ipsilateral hypothalamus, brainstem tegmentum, bilateral precunei, and lateral parietal lobes, mainly with early time-shift models, and the contralateral dominant bilateral cerebellum (contralateral dominant) with later time-shift models. Deactivations included the ipsilateral cuneus, bilateral caudate nuclei, thalamus, hippocampi, precunei, and lateral parietal lobes (Usami et al., 2016).

means of comprehensive study such as video-EEG monitoring (VEEG), neuropsychological test, FDG-PET and MRI, we have revealed the smoldering nature of the autoimmune limbic encephalitis. We have been utilizing multidisciplinary approach (VEEG, FDG-PET and 3T MRI) for offering tailor-made immunotherapy for patients with autoimmune epilepsy. Additionally, we reported that autoimmune mechanism could underlie in an epileptic patient with atypical clinical picture (triple pathology in a patient with parietal tumor) (Matsumoto et al., 2015). Now we attempt to make diagnostic criteria of autoimmune epilepsy, based on both clinical picture and various examinations (Sakamoto et al., 57th annual meeting of Japan Society of Neurology).

We started conducting collaborative researches in epilepsy by means of iPS cells with Prof. Haruhisa Inoue (Center for iPS Cell Research and Application: CiRA).

c) Advanced EEG analysis

EEG now has a history of more than 80 years for evaluation of brain functions and diagnosis of brain diseases, ranging from brain death, coma to epilepsy. Medico-engineering collaboration between Prof. Shibasaki's group (Kyoto University School of Medicine, Neurology, Human Brain Research Center) and Prof. Nakamura's group (Saga University, Faculty of Science and Engineering) has been done to develop the automatic EEG interpretation system, and it is currently under clinical investigation. Now we plan to establish a remote EEG viewing system in collaboration with the department of Neurology of Kobe City Medical Center General Hospital and Takeda General Hospital, and Nihon Kohden, aiming at improvement in quality about clinical practice of epilepsy and a close intermural link. In addition, we started a collaborative study with National Cerebral and Cardiovascular Center in order to establish the diagnostic criteria and treatment for the post-stroke epilepsy patients.

2) Mapping higher functions/network and elucidating its functional alteration under pathological condition

In epilepsy surgery, it is important to map cortical functions to preserve eloquent cortices in addition to the localization of the epileptic focus. Therefore, we need to perform comprehensive 'system mapping' to help neurosurgeons to make strategy of surgery for individual patients. We have made vigorous attempts at developing various techniques for mapping higher cortical functions (e.g., language, motor control etc.) and their network for clinical application.

Functional neuroimaging tells us if specific brain regions are active during certain tasks, but activation by itself does not demonstrate the necessity of those areas. In contrast, electrical cortical stimulation, a gold standard method since mid-20th century, can delineate the cortex responsible for a particular task by making functional impairment. The functional interference is temporary (~ 5 s), discretely focal ($\sim 1 \text{ cm}^2$), and in sharp contrast to chronic stroke lesions that are relatively large and usually associated with cortical plastic compensation.

However, high frequency electrical stimulation often results in afterdischarges that delay functional mapping and harbor a risk of seizure induction. Recent technical advances have enabled us to record the cortical activities relevant to higher cortical functions with wideband EEG technology - from infraslow to high gamma activities. In our institute, in addition to the gold standard method of high frequency electrical stimulation, we perform comprehensive mapping of higher cortical functions by recording epicortical infraslow and high frequency oscillation/activity during motor or cognitive tasks. Recently, we investigated the role of the basal temporal language area (BTLA) using a comprehensive approach. Discrete local field potentials were observed in BTLA while participants completed semantic tasks. Electrical stimulation (lower stimulus intensity & duration) to the area evoked semantic paraphasia and prolonged reaction time selectively during the semantic tasks. The study demonstrated that the BTLA is crucial for semantic processing especially among language functions (Shimotake et al. 2015). In some researchers, it has been considered that the basal anterior temporal lobe is the 'visual' language area. To identify whether this area has visual or semantic function, we, for the first time, applied the representational similarity analysis (RSA), one of the novel decoding methods, to ECoG data (epicortical event related potentials recorded during picture naming task). We could demonstrate that ventral anterior temporal lobe selectively coded semantic processing rather than the visual

or phonological processing (Fig. 3) (Chen, Shimotake et al. 2016). Furthermore, high frequency electrical stimulation to the area and its adjacent areas sometimes induced mirth or laughter, suggesting that this area is also related to semantic processing of humor (Yamao et al., 2016).

We incorporate cortico-cortical evoked potentials (CCEPs) to probe inter-areal functional connectivity in order to perform 'system mapping'. CCEP is an *in vivo* electrical tract tracing method developed in Cleveland and Kyoto (Matsumoto et al., 2004). Single electrical stimulation (1 Hz) is applied to a part of the cortices and cortical evoked potentials, i.e., CCEPs, were recorded from adjacent and remote cortical activities through cortico-cortical connections. With this CCEP method, we can probe cortico-cortical networks involved in functional brain systems and seizure network in each individual patient. Furthermore, by gathering data of cortical functions and networks from many patients in various physiological and pathological states and analyzing them as a group, we attempt to feedback this valuable information into the system neuroscience by providing functional/connectivity references for non-invasive researches.

We can also use induced evoked potential (CCEP) and high frequency activities (HFA) as a dynamic marker of cortical connectivity and excitability since it can "snapshot" the connectivity and excitability within 20-30 trials (< 1 min). We applied this method to awake and sleep cycle and found that cortical connectivity and excitability are different between wakefulness and non-REM (rapid eye movement) sleep, with REM sleep being the intermediate state among them. Increase of neuronal excitability, as reflected in induced HFA, was immediately followed by intense inhibition during non-REM sleep. It was only the frontal lobe that showed rebounded hypersynchrony after intense inhibition during non-REM sleep. These sleep-induced dynamic changes or properties would underlie unconsciousness during sleep and frequent nocturnal seizures in frontal lobe epilepsy.

HFAs consist of physiological HFAs related to various brain functions and pathological (epileptic) HFAs or high frequency oscillations (HFOs). Factors regulating HFA frequency have not been well elucidated in humans. We focused on the primary somatosensory cortex (SI) since physiological HFAs overriding somatosensory evoked potentials have




Figure 3. Representational similarity analysis (RSA) applied to the cortical event related potential

We used representational similarity analysis (RSA) to the cortical event related evoked potential during picture naming task performed in ten patients with subdural electrodes covering ventral and lateral anterior temporal regions.

We compared the obtained neural activity with the pre-defined theoretical semantic, visual and phonological models. Neural activity selectively coded the semantic representation in the ventral anterior temporal subregion. (Modified from Chen, Shimotake et al., 2016 with permission)

been extensively studied in humans. We, in the SI, compared HFAs induced by median nerve stimulation with those induced by electrical stimulation (HFAs overriding CCEPs) of the cortex connecting to SI. We clarified that different propagation modes and/or different terminal layers seemed to determine HFA frequency, and that since HFAs overriding CCEPs and HFA induced during various brain functions share a similar broadband profile of the power spectrum, cortico-cortical horizontal propagation seems to represent common mode of neural transmission for processing these functions (Kobayashi et al., 2015).

We have demonstrated the central mechanisms and functional alteration under pathological condition relevant to i) the motor control (negative motor phenomena, praxis, reaching, conflict processing and response inhibition), ii) language (dorsal and ventral language networks with emphasis on semantic cognition) and iii) visual functions (retinotopic mapping by functional MRI), combined with non-invasive evaluation (functional MRI, diffusion tractography, MEG, neuropsychology). Additionally, we are now tackling with decoding of complex neural signals during various tasks in cooperation with seasoned researchers in and out of the Kyoto University (Graduate School of Informatics, Dr. Satoshi Tsujimoto; Advanced Telecommunications Research Institute International, Dr. Rieko Osu; School of Psychological Sciences, the University of Manchester, Prof. Matthew Lambon-Ralph). Finally, we are pleased to announce that our group (A03 “the Direct Recording of Human Neural Oscillations”, Prof. Ikeda as PI) engages in the national research group of “Neuro-Oscillology” which is funded by Grant-in-Aid for Scientific Research on Innovative Areas from the Ministry of Education, Culture, Sports, Science and Technology (MEXT). We also have taken part in the ‘Embodied-Brain Systems Science’ funded by another grant from MEXT (A03-4, Dr. Riki Matsumoto as PI), in order to reveal the motor control and the mechanism of body representations especially in the fronto-parietal network.

3) Pathogenesis of movement disorders and its treatment

We have investigated movement disorders, mainly myoclonus and myoclonus epilepsy, by way of epidemiological, genetic and electrophysiological methods. BAFME (benign adult onset familial myoclonus epilepsy) has been investigated mainly in Japan and European countries for 20 years. The clinical pictures are as follows: i) adult onset, ii) autosomal dominant (unknown causative gene), iii) cortical (myoclonic) tremor (tremulous myoclonus), iv) infrequent generalized seizure, v) cortical reflex myoclonus disclosed by electrophysiological study. We have also been studying BAFME since it was first reported in 1990. As its name suggests, BAFME was considered to present no progression




Figure 4. Linear regression analysis between age at the time of EEG recording and frequency of PDR in 19 BAFME patients and 38 age-matched control subjects. A simple regression curve for BAFME patients is represented by the black line, and for age-matched control subjects by the grey dotted line. Open circles indicate BAFME patients and crosses indicate age-matched control subjects. There was a non-significant tendency for a slowing of PDR with age in both BAFME and age-matched control subjects. Comparison of the gradients corresponding to PDR frequency relative to age between the two groups revealed no significant difference.

and good prognosis. However, cortical myoclonic tremor has been proved to worsen with aging. Recently, we demonstrated slow progression of the disease, based on the electrophysiological evidence. Namely, the amplitude of somatosensory evoked potential, reflecting the cortical excitability in the primary sensori-motor cortices, more exaggerated with aging in BAFME patients than normal volunteers. We also demonstrated clinical anticipation in BAFME, in which the onset of generalized seizure and cortical (myoclonic) tremor became earlier in the next generation. The anticipation in BAFME was more apparent in patients with maternal transmission. By comparing the EEG posterior dominant rhythms (PDRs) between patients with BAFME and age-matched control subjects, we showed mild diffuse encephalopathy in BAFME. By means of cortico-muscular coherence, we clarified that hyperexcitability of primary sensorimotor cortex and the subcortical structures such as basal ganglia and cerebellum would be involved in the generator mechanisms of cortical tremor in BAFME (in preparation). These findings would be helpful to search a causative gene of BAFME. In addition, the nationwide questionnaire for neurologists and epileptologists in Japan revealed that BAFME patients were found diffusely without regional accumulation. The further survey based on the detailed clinical information of about 100 BAFME patients showed that the clinical core symptoms were rather homogeneous for BAFME in Japan (in preparation).

Unverricht-Lundborg disease (ULD) is the most common form of progressive myoclonus epilepsy syndrome (PME) in the world, but mainly reported from Baltic and Mediterranean region. The symptoms consist of epileptic seizure, myoclonus, ataxia and cognitive impairment and are gradually deteriorated. We reported ULD cases in Japan, in some of which the development of symptoms and increase of SEP amplitude became very subtle for long-term follow-up. Similar to BAFME, PME including ULD also presented no regional clustering in Japan.

Besides epilepsy patients, we reported a case of elderly woman with exaggerated startle reflex and unconsciousness drop attack including the analysis of electromyography polygraph.

Collaborators

We have been collaborating closely with the Departments that officially support our department. Other collaborators are listed below.

[Overseas]

Dr. Stéphanie Baulac, Ph.D.

Affiliation: Institut du Cerveau et de la Moelle épinière (ICM), Epilepsy Unit

Position: Research Director

Dr. Christophe Bernard, Ph.D.

Affiliation: INS - Institut de Neurosciences des Systèmes, UMR INSERM 1106, Aix-Marseille Université

Position: Team leader

Prof. Marco Catani, M.D., Ph.D.

Affiliation: Natbrain lab, Department of Forensic and Neurodevelopmental Sciences, Institute of Psychiatry, King's College London

Position: Head of the Natbrainlab, Clinical Senior Lecturer and Honorary Consultant Psychiatrist

Prof. Nathan Earl Crone, M.D.

Affiliation: Department of Neurology, Johns Hopkins University School of Medicine

Position: Professor

Prof. Matthew A. Lambon-Ralph, FRCSLT (hons), FBPsS

Affiliation: School of Psychological Sciences, University of Manchester

Position: Professor of Cognitive Neuroscience & Associate Vice-President (Research)

Dr. Dileep R. Nair, M.D.

Affiliation: Epilepsy Center, Cleveland Clinic

Position: The Section Head of Adult Epilepsy and Director of Intraoperative Neurophysiologic monitoring

Prof. Angela Vincent, Ph.D.

Affiliation: University of Oxford

Position: Emeritus professor

Dr. Marco de Curtis, M.D.

Affiliation: Fondazione IRCCS Istituto Neurologico Carlo Besta

Position: Head of Epileptology and Experimental Neurophysiology Unit, Head of Pre-clinical Neuroscience Laboratories

[Domestic]

Dr. Koji Iida, M.D., Ph.D.

Affiliation: Department of Neurosurgery, Hiroshima University Hospital

Position: Associate Professor

Dr. Yushi Inoue, M.D., Ph.D.

Affiliation: Shizuoka Institute of Epilepsy and Neurological Disorders, National Epilepsy Center, Department of Clinical Research

Position: Hospital director

Prof. Shigeki Kameyama, M.D., Ph.D.

Affiliation: Nishi-Niigata Chuo National Hospital

Position: Honorary hospital director

Prof. Amami Kato, M.D., Ph.D.

Affiliation: Department of Neurosurgery, Kinki University Hospital

Position: Professor

Prof. Masatoshi Nakamura, Ph.D.

Affiliation: Research Institute of Systems Control, Institute for Advanced Research and Education, Saga University

Position: Emeritus professor

Prof. Shigeto Nishida, Ph.D.

Affiliation: Department of Information and Communication Engineering, Faculty of Information Engineering, Fukuoka Institute of Technology

Position: Professor

Dr. Teiichi Onuma, M.D., Ph.D.

Affiliation: Musashino Kokubunji Clinic

Position: Honorary hospital director

Dr. Rieko Osu, Ph.D.

Affiliation: Department of Motor Control and Rehabilitation, ATR Computational neuroscience Labs

Position: Department Head

Dr. Satoru Saito, Ph.D.

Affiliation: Division of Cognitive Psychology in Education, Kyoto University Graduate School of Education

Position: Professor

Prof. Yoshio Sakurai, Ph.D.

Affiliation: Laboratory of Neural Information, Graduate School of Brain Science, Doshisha University

Position: Professor

Dr. Hiroshi Shibasaki, M.D., Ph.D.

Affiliation: Kyoto University

Position: Professor Emeritus

Dr. Takenao Sugi, Ph.D.

Affiliation: Institute of Ocean Energy, Saga University

Position: Associate professor

Prof. Shoji Tsuji, M.D., Ph.D.

Affiliation: Department of Neurology, The University of Tokyo Hospital

Position: Professor

Dr. Satoshi Tsujimoto, Ph.D.

Affiliation: Department of Intelligence Science and Technology, Graduate School of Informatics, Kyoto University

Position: Associate professor

Dr. Hiroki Yamamoto, Ph.D.

Affiliation: Graduate School of Human and Environmental Studies, Kyoto University

Position: Assistant professor

Dr. Ikuko Yano, Ph.D.

Affiliation: Department of Pharmacokinetics and Pharmaceutics, Kobe University Graduate School of Medicine

Position: Associate professor

(Listed in the alphabetical order of their family names)

Direct recording of the neural oscillation in human brain

(Grant-in-Aid for Scientific Research on Innovative Areas from the Ministry of Education, Culture, Sports, Science and Technology, Japan: Non-linear Neuro-oscillology: Towards Integrative Understanding of Human Nature)

Prof. Ichiro Tsuda, Ph.D.

Affiliation: Department of Mathematics, Faculty of Science, Hokkaido University

Position: Professor

Dr. Keiichi Kitajo, Ph.D.

Affiliation: Rhythm-based Brain Information Processing Unit, RIKEN Brain Science Institute

Position: Unit Leader

Prof. Katsunori Kitano, Ph.D.

Affiliation: Department of Human and Computer Intelligence, Ritsumeikan University

Position: Professor

Prof. Toshio Aoyagi, Ph.D.

Affiliation: Department of Applied Analysis and Complex Dynamical Systems, Kyoto University Graduate School of Informatics

Position: Professor

Prof. Tatsuya Mima, M.D., Ph.D.

Affiliation: The Graduate School of Core Ethics and Frontier Sciences, Ritsumeikan University

Position: Professor

Dr. Hiroaki Wagatsuma, Ph.D.

Affiliation: Graduate School of Life Science and Systems Engineering, Kyushu Institute of Technology

Position: Associate Professor

Prof. Katsuhiro Kobayashi, M.D., Ph.D.

Affiliation: Department of Child Neurology, Okayama University Graduate School of Medicine

Position: Professor

Prof. Takashi Nagamine, M.D., Ph.D.

Affiliation: Department of Systems Neuroscience, Sapporo Medical University School of Medicine

Position: Professor

Neural basis of human body representation: a direct electrocorticographic recording and stimulation study.

(Grant-in-Aid for Scientific Research on Innovative Areas from the Ministry of Education, Culture, Sports, Science and Technology, Japan: Understanding brain plasticity on body representations to promote their adaptive functions ("Embodied-Brain")

Dr. Eiichi Naito, Ph.D.

Affiliation: Center for Information and Neural Networks (CiNet), National Institute of Information and Communications Technology (NICT)

Position: Research Manager

Prof. Hiroshi Imamizu, Ph.D.

Affiliation: Department of Intelligence Science and Technology, Graduate School of Informatics, Kyoto University

Position: Professor

Dr. Takaki Maeda, M.D., Ph.D.

Affiliation: Department of Neuropsychiatry, Keio University School of Medicine

Position: Senior Lecturer

The elucidation of glial function in intractable epilepsy and standardization of clinical practice guidelines.

(Japan Agency for Medical Research and Development (AMED): Practical Research Project for Rare Diseases)

Prof. Taketoshi Maehara, M.D., Ph.D.

Affiliation: Department of Neurosurgery, Tokyo Medical and Dental University

Position: Professor

Prof. Akiyoshi Kakita, M.D., Ph.D.

Affiliation: Department of Pathology, Brain Research Institute, Niigata University

Position: Professor

Prof. Yukihiro Ohno, Ph.D.

Affiliation: Laboratory of Pharmacology, Osaka University of Pharmaceutical Sciences

Position: Professor

The establishment of diagnostic criteria and treatment for post-stroke epilepsy patients.

(AMED: Practical Research Project for Life-Style related Diseases including Cardiovascular Diseases and Diabetes Mellitus)

Dr. Masafumi Ihara, M.D., Ph.D.

Affiliation: Department of Stroke and Cerebrovascular Diseases, National Cerebral and Cardiovascular Center

Position: Director, Cerebrovascular Medicine and Neurology

Clinical activities

1) Outpatient Epilepsy Clinic. Promoting cooperation between hospitals and clinics for epilepsy care

As a team of specialists, we have made full efforts to provide the best care to patients suffering from epilepsy or movement disorders. Until recently, epilepsy has been recognized as a childhood-onset disease. However, with the advent of a superaging society, epilepsy that develops in the middle-aged or elderly has become a current problem in Japan. In addition, the number of the hospitals and physicians that can offer the epilepsy care is not adequate. Moreover, it is unclear which department, neurology, neurosurgery, or psychiatry, is in charge for the adult epilepsy service. In order to offer the optimal epilepsy care, it is very important to establish the cooperation model among general physicians and epilepsy specialists for epilepsy care like that in European and North American countries. As a tertiary care institute for epilepsy in Kyoto, we have led cooperation among primary, secondary and tertiary facilities in the Kinki district (esp. in Kyoto-Shiga region) to provide a comprehensive epilepsy service with a dedicated team of neurologists, neurosurgeons, pediatricians and psychiatrists.

In the fiscal year 2015, we saw 1008 outpatients. 299 patients were newly consulted from other hospitals and clinics in the Kinki district. We promoted hospital-clinic cooperation by returning the referral patients to their local clinics and hospitals.

2) Inpatient evaluation and treatment for epilepsy (including video-EEG monitoring)

Since 1991, we have been running the epilepsy monitoring unit (EMU) in the Neurology Ward for evaluation of patients with epilepsy. We now have two dedicated rooms for EMU, equipped with the digital video-EEG system.

By capturing seizures with simultaneous video and EEG recording, we can perform

- i) An accurate diagnosis of epilepsy: To determine whether the seizure is epileptic or non-epileptic, including movement or psychogenic disorders,
- ii) Identification of epileptic focus: To locate the epileptic focus for epilepsy surgery in patients with medically intractable epilepsy.

In the fiscal year 2015, we examined 40 patients in the EMU (subdural/depth electrode implantation: 3, presurgical evaluation: 15, evaluation of limbic encephalitis: 5, diagnosis of epilepsy: 17).

In addition, we provide patients with multidisciplinary studies for comprehensive evaluation, such as 3 tesla MRI, routine EEG, FDG-PET/SPECT, MEG and neuropsychological testing. Routine EEGs were performed in 1231 patients (including 869 outpatients) in this fiscal year.

3) Epilepsy Surgery

We have established an epilepsy surgery program with close collaboration with the Department of Neurosurgery since 1991. Since the first epilepsy surgery in 1992, we have performed more than 190 epilepsy surgeries, with the majority of patients having seizure freedom or substantial decrease leading to better QOL. We provide each patient with the individually tailored surgery plan by incorporating the findings of the multimodal studies (see below) as well as the Wada test. The patients may proceed to the invasive presurgical evaluation with intracranial electrodes (subdural and/or depth electrodes) when the epileptic focus cannot be precisely localized (such as in cases with non-lesional MRI) or the focus is located at or around the functionally important areas such as motor or language cortices. In such cases, the patients undergo the first surgery for implantation of intracranial electrodes. After electrode implantation, the patients are evaluated for the epileptic focus (by recording seizures) and the functional cortical areas (by incorporating the state-of-art mapping techniques) for 1-2 weeks. Then, the patients undergo the second surgery for resection of the epileptic focus. The patients may undergo the awake brain surgery, where the patients wake up from anesthesia if necessary. Awake surgery has the advantage to evaluate the brain functions such as motor and language during resection and monitor the 'natural' epileptic spikes without any influence from anesthetics. In the fiscal 2015, our team performed epilepsy surgery in 8 patients (3 with chronic intracranial electrode implantation) and awake brain surgery about 24 patients (including non-epilepsy cases).

4) Examinations for epilepsy

As the tertiary care epilepsy facility, we provide patients with the state-of-arts studies for the evaluation of epilepsy. As the comprehensive epilepsy program in the national university hospital, we incorporate the leading techniques as clinical research studies (IRB approved) for the optimal presurgical evaluations.

- Electroencephalography (EEG)
 - Magnetoencephalography (MEG)
 - FDG-PET (18F-fluorodeoxyglucose positron emission tomography)
SPECT (Single photon emission computed tomography) including ictal SPECT
 - 3 tesla MRI
 - functional MRI (fMRI)
 - EEG-fMRI (simultaneous EEG and functional MRI recording)
 - Neuropsychological testing (WAIS-III, WMS-R, WAB, semantic batteries and Kanji/Kana related tasks)
 - invasive EEG monitoring with intracranial electrodes
- Recently, autoimmune epilepsy is regarded as one of the important cause of epilepsy. Following tests are diagnostic for autoimmune epilepsy.
- Cerebrospinal fluid / serum antibody test

5) Development of novel treatments for epilepsy

- i) Interventional Neurophysiology: Recently, neurophysiology has been highlighted for its application to treatment of various neurological diseases. In our hospital, we apply a novel interventional neurophysiology method, neurofeedback treatment, to medically intractable patients in whom epilepsy surgery is not applicable. Patients train themselves to control the brain activity (by adjusting slow EEG potentials) to suppress epileptic seizure activity. Our preliminary study shows a good efficacy as comparable to that for the Vagus Nerve Stimulation.
- ii) Promoting the clinical trials for new anti-epileptic drugs.

6) Diagnosis and treatment for movement disorders

It is also our mission to provide the optimal care for patients with movement disorders. We provide precise diagnosis using advanced diagnostic tools for better treatment of movement disorders such as tremor, myoclonus, dystonia and other involuntary movements.

The pathophysiology of movement disorders, however, is not fully understood. We have been investigating their pathophysiology and treatment in close collaboration with the Department of Neurology and Human Brain Research Center (HBRC).

7) Simulation training of brain death determination

Since Organ Transplant Law went into force in 1997, we, in close collaboration with the affiliated departments, have been regularly practicing the course 'Simulation-based training in brain death determination'. In this course, we simulate the management about how and what to do when the donor is found and until organs are taken. The training is highly practical for those in charge of brain death determination in our hospital. In the fiscal 2015, we participated in the first case of brain death determination of Kyoto University Hospital in close collaboration with many affiliated departments.

京都大学大学院医学研究科 / 京都大学医学部附属病院神経内科 寄附講座
てんかん・運動異常生理学講座
Department of Epilepsy, Movement Disorders and Physiology

HOME | サイトマップ English 検索 

HOME

- ご挨拶
- 診療内容
- メンバー
- 研究
- 業績
- 教育
- 年次報告書 Annual Report
- リンク

**京都大学医学部附属病院
てんかん外来のご案内** 

京滋てんかん病診連携 

新時代のてんかんの診療を実現させる Education Clinical Care Research

てんかん・運動異常生理学講座の御紹介

本講座では、以下を設置理由および目的として、京都大学医学研究科に寄附講座として、平成25年8月1日に設立されました。本学の臨床神経学講座(高橋良輔教授)より支援講座としての御支援をうけながら、以下を目指します。

- 集学的立場から、てんかんおよび運動異常症の病態解明と治療の開発、高度先進医療の推進、実践医療としての確立と普及を目指す。同時に、将来の本分野の担い手となる専門医と臨床研究者の養成と教育機会を国内外に広く提供します。
- 臨床てんかん学の病態と治療と常に表裏一体関係である生理学の研究と臨床応用の発展を、大学病院の立場から推進します。

診療 研究 教育


お知らせ

Video EEG monitoring room



デジタル脳波判読の遠隔診断

データ通信での判読、患者負担なし、正確な診断と迅速な対応



(reprint permission from Nihon Koden)

EEG conference



Annual course 'Simulation-based training in brain death determination' in Kyoto University Hospital



Educational activities

1) Offering the optimal education and research to Japanese and foreign physicians

■ EEG/Epilepsy fellowship

With great support by the Department of Neurology, we have set up the EEG/Epilepsy fellowship for training young neurologists, neurosurgeons, pediatricians, and psychiatrists. Four adult neurologists, one pediatric neurologist, and one neurosurgeon have already completed this fellowship. Our education covers various fields of epileptology with a focus on clinical neurophysiology. We plan to welcome foreign young doctors for fellowship training as well.

Contents of the fellowship program are listed as follows;

- 1) Training of routine EEG reading (emergency EEG as well)
- 2) Analysis of the long-term video-EEG monitoring for diagnosis and presurgical evaluation
- 3) Clinical practice of adult epilepsy
- 4) Training of medical treatment with anti-epileptic drugs

Graduates of EEG/Epilepsy fellowship

Reiko Tsuda (from June 2011 to August 2011)

Daiki Fujii (from September 2014 to November 2014)

Takeshi Inoue (from April 2013 to March 2016)

Hajime Yoshimura (from July 2015 to September 2015)

Tsuyoshi Tsukada (from October 2015 to March 2016)

Current Trainee of EEG/Epilepsy fellowship

Daiki Fujii (from February 2016 -)

Masayuki Honda (from April 2016 -)

■ Intramural, multidisciplinary monthly case conference

In cooperation of the Departments of Neurology, Neurosurgery, Pediatrics, Diagnostic Radiology, Psychiatry, Rehabilitation, and Clinical Laboratory Medicine, and Human Brain Research Center, we have been holding the intramural, multidisciplinary monthly case conference for more than a decade. In the conference, we discuss the diagnosis and surgical indication of epilepsy patients for comprehensive epilepsy practice as a tertiary epilepsy special facility. The numbers of participants and the discussion cases are getting larger. As a training facility certified by Japan Epilepsy Society (JES), this conference is open for doctors outside the hospital to discuss their problem case or to obtain the credit to apply board examination of JES-certified epileptologist.

■ EEG conferences and so on

For our graduate students and EEG/Epilepsy fellows, we have been offering multifaceted educational and research trainings, such as EEG reading skills in EEG conferences twice a week, seeing outpatients and inpatients with staffs, and epilepsy/clinical neurophysiology researches. One EEG conference and research conference are held in English for training skills in English presentation. The other conference is held in Japanese and open for the in-hospital technicians and out-hospital doctors for providing them with training opportunities for the practical basic EEG reading skills (about 30-40 participants).

■ Specialist training

In the fiscal year 2015, our department produced one board-certified epileptologist (JES).

■ Extramural workshops

Regarding educational activities outside the institute, as the secretary office in general, we have organized the district EEG & EMG teaching course for the young doctors and technicians in Kansai (Kansai EEG & EMG workshop) every year since 2008. In 2015, we newly founded and organized the EEG seminar advanced course for the purpose of acquisition of specialized knowledge and reading skills in clinical EEG sponsored by the Japanese Society of Clinical Neurophysiology. We also have provided educational activities by complying the request of lectures nationwide (please refer to the achievements for details). Both staffs regularly teach EEG reading and epileptology at the affiliated hospitals.

■ EEG/Epilepsy lecture series

We have been holding the intensive lecture series of the basics of EEG and epileptology for EEG/Epilepsy fellows and young doctors.

2) Offering medical staffs' education for caring of epilepsy patients

In the Kyoto University Hospital, we have offered education for epilepsy and related disorders to doctors and medical staffs. For the medical staffs in the Neurology clinic and ward, we hold comprehensive monthly lectures about pathophysiology of epilepsy, seizure semiology, and medical care of patients living with epilepsy.

3) Providing patients, family and society with valuable information

We have responded to the request by the patients, family, and society in cooperation with Japan Epilepsy Association. For example, we have joined the lectures sponsored by Japan Epilepsy Association for the public, and also the continuing medical education lectures for physicians by Japan Medical Association.

Interdisciplinary monthly case conference



Interdisciplinary monthly case conference



EEG/Epilepsy lecture series



EEG reading room (Department of Neurology)



Recruitment of EEG/Epilepsy fellowship



京都大学神経内科 EEG/Epilepsy Fellowship 募集

1. 対象 卒後4年目以降の段階の神経内科、脳外科、小児科、精神科等の若手医師
2. 専門研修内容
 - i. 幅広い脳波判読の研修と経験
 - ii. 長時間ビデオ脳波モニターの解析
 - iii. てんかんの診療
 - iv. 各種抗てんかん薬の臨床研修
3. 期間 単年度単位で1年間（より短期研修も希望に応じて可能）
4. 処遇 非常勤医師待遇

少しでも興味がある方は、一度是非ご連絡ください

連絡先：京都大学神経内科事務室、担当：松本理器、池田昭夫
tel: 075-751-3771, fax: 075-751-3265, email: neurofc@kuhp.kyoto-u.ac.jp

Research grants obtained from extramural sources & awards

The ministry of Education, Culture, Sports, Science and Technology of Japan Grant-in-Aids for Scientific Research (KAKENHI)

Fiscal years 2015 – 2019

Grant-in-Aid for Scientific Research on Innovative Areas (Non-linear Oscillology)

Principal investigator: Akio Ikeda

Subject number: 15H05874

Fiscal years 2015 – 2016

Grant-in-Aid for Scientific Research on Innovative Areas (Embodied Brain Systems Science)

Principal investigator: Riki Matsumoto

Subject number: 15H01664

Fiscal years 2014 – 2016

Grant-in-Aid for Scientific Research (B)

Principal investigator: Akio Ikeda

Subject number: 26293209

Fiscal years 2014 – 2017

Grant-in-Aid for Scientific Research (B)

Principal investigator: Riki Matsumoto

Subject number: 26282218

Health Labour Sciences Research Grant

Fiscal years 2014 – 2016

Principal investigator: Yushi Inoue

Co-investigator: Akio Ikeda

Subject number: H26 - 難治等 - 一般 - 051

Japan Agency for Medical Research and Development (AMED)

Fiscal years 2015 – 2017

Co-investigator: Akio Ikeda

Subject number: 15ek0109120h0001

Fiscal years 2016 – 2017

Co-investigator: Akio Ikeda

Subject number: 16ek0210057h0001

Others

The Japan Epilepsy Research Foundation Research Grant

Fiscal years 2014 – 2016

Principal investigator: Riki Matsumoto

The Japan Epilepsy Research Foundation Research Grant

Fiscal years 2014 – 2016

Principal investigator: Tomoyuki Fumuro

Awards

Akio Ikeda: Masakazu Seino Memorial Lecture

The 11th Asian & Oceanian Epilepsy Congress (AOEC), Hong Kong, May 2016

Riki Matsumoto, Takeharu Kunieda: The Japan Epilepsy Research Foundation Research Award 2016

IV Society-academia collaboration

As society-academia collaboration, both collaborative study and teaching seminar were conducted. All activities were conducted with the companies under the regulation of Kyoto University Graduate School of Medicine and University Hospital and under the regulation of Japanese law. Therefore, the detailed English version was not made, and please refer to the Japanese part of this report in details.

Only the name of company and the content of collaboration were listed as follows.

1) Collaborative study

GlaxoSmithKline K.K.

Clinical study of drug exchange from Valproic acid to Lamotrigine

NIHON KOHDEN CORPORATION

- 1) Analysis system of wide-band EEG in clinical EEG
- 2) Clinical and technical establishment of remote reading system of digital EEG in Japan
- 3) Optimal condition of high frequency cortical stimulation in cortical mapping

UCB Japan Co., Ltd.

Phase III clinical trial of Lacosamide

2) Teaching seminars

UCB Japan Co., Ltd.

Clinical epilepsy seminar in Kinki district area: a whole day course in 2014 and 2015



業績報告 Publications and Congress Presentations

誌上発表 Publications

原著 Original articles

〈英文 English articles〉

1. Usami K, Matsumoto R, Sawamoto N, Murakami H, Inouchi M, Fumuro T, Shimotake A, Kato T, Mima T, Shirozu H, Masuda H, Fukuyama H, Takahashi R, Kameyama S, Ikeda A: Epileptic network of hypothalamic hamartoma: An EEG-fMRI study. *Epilepsy Res* 2016, 125: 1-9.
2. Chen Y, Shimotake A, Matsumoto R, Kunieda T, Kikuchi T, Miyamoto S, Fukuyama H, Takahashi R, Ikeda A, Lambon Ralph MA: The 'when' and 'where' of semantic coding in the anterior temporal lobe: temporal representational similarity analysis of electrocorticogram data. *Cortex* 2016, 79: 1-13.
3. Imamura H, Matsumoto R, Takaya S, Nakagawa T, Shimotake A, Kikuchi T, Sawamoto N, Kunieda T, Mikuni N, Miyamoto S, Fukuyama H, Takahashi R, Ikeda A: Network specific change in white matter integrity in mesial temporal lobe epilepsy. *Epilepsy Res* 2016, 120: 65-72.
4. Ito S, Yano I, Hashi S, Tsuda M, Sugimoto M, Yonezawa A, Ikeda A, Matsubara K: Population Pharmacokinetic Modeling of Levetiracetam in Pediatric and Adult Patients With Epilepsy by Using Routinely Monitored Data. *Ther Drug Monit* 2016, 38: 371-378.
5. Yamao Y, Kunieda K, Matsumoto R: Reply to commentary on "neural correlates of mirth and laughter: a direct electrical cortical stimulation study". *Cortex* 2016, 75: 244-246.
6. Hitomi T, Kobayashi K, Sakurai T, Ueda S, Jingami N, Kanazawa K, Matsumoto R, Takahashi R, Ikeda A: Benign adult familial myoclonus epilepsy is a progressive disorder: no longer idiopathic generalized epilepsy. *Epileptic Disord* 2016, 18: 67-72.
7. Neshige S, Kobayashi K, Shimotake A, Iemura T, Matsumoto R, Nishinaka K, Matsumoto M, Takahashi R, Ikeda A: An elderly woman with exaggerated startle reflex and unconscious drop attack. *Neurology and Clinical Neuroscience* 2016, in press. DOI : 10.1111/ncn.12055.
8. Uneno Y, Yokoyama A, Nishikawa Y, Funakoshi T, Ozaki Y, Aoyama I, Yamaguchi D, Morita S, Mori Y, Kanai M, Kinoshita H, Inoue T, Sawamoto N, Matsumoto R, Matsumoto S, Muto M: Paraneoplastic limbic encephalitis in human epidermal growth factor receptor-2-positive gastric cancer patient treated with trastuzumab-combined chemotherapy: case report and literature review. *Internal Medicine* 2016, in press.
9. Fukuma K, Ihara M, Miyashita K, Motoyama R, Tanaka T, Kajimoto K, Ikeda A, Nagatsuka K: Right parietal source in Mahjong-induced seizure: a system epilepsy of focal origin. *Clinical Case Reports* 2016, in press.
10. Ritaccio A, Matsumoto R, Morrell M, Kamada K, Koubeissi M, Poeppel D, Lachaux JP, Yanagisawa Y, Hirata M, Guger C, Schalk G: Proceedings of the Seventh International Workshop on Advances in Electrocorticography. *Epilepsy Behav* 2015, 51: 312-320.
11. Ikeda A: Commentary: We shall not lose past medical documents. *Epilepsia* 2015, 56: 1712-1713.
12. Usami K, Matsumoto R, Kobayashi K, Hitomi T, Shimotake A, Kikuchi T, Matsuhashi M, Kunieda T, Mikuni N, Miyamoto S, Fukuyama H, Takahashi R, Ikeda A: Sleep modulates cortical connectivity and excitability in humans: Direct evidence from neural activity induced by single-pulse electrical stimulation. *Hum Brain Mapp* 2015, 36: 4714-29.

13. Yamao Y, Matsumoto R, Kunieda K, Arakawa Y, Kikuchi T, Shibata S, Ikeda A, Fukuyama H, Miyamoto S: A possible variant of negative motor seizure arising from the supplementary negative motor area. *Clin Neurol Neurosurg* 2015, 134: 26-129.
14. Shimotake A, Matsumoto R, Ueno T, Kunieda T, Saito S, Hoffman P, Kikuchi T, Fukuyama H, Miyamoto S, Takahashi R, Ikeda A, Lambon Ralph MA: Direct Exploration of the Role of the Ventral Anterior Temporal Lobe in Semantic Memory: Cortical Stimulation and Local Field Potential Evidence From Subdural Grid Electrodes. *Cereb Cortex* 2015, 25: 3802-3817.
15. Yamao Y, Matsumoto R, Kunieda T, Shibata S, Shimotake A, Kikuchi T, Satow T, Mikuni N, Fukuyama H, Ikeda A, Miyamoto S: Neural correlates of mirth and laughter: a direct electrical cortical stimulation study. *Cortex* 2015, 66: 134-140.
16. Kobayashi K, Matsumoto R, Matsuhashi M, Usami K, Shimotake A, Kunieda T, Kikuchi T, Mikuni N, Miyamoto S, Fukuyama H, Takahashi R, Ikeda A: Different Mode of Afferents Determines the Frequency Range of High Frequency Activities in the Human Brain: Direct Electrocorticographic Comparison between Peripheral Nerve and Direct Cortical Stimulation. *PLoS One* 2015, 10: e0130461.
17. Inoue Y, Yagi K, Ikeda A, Sasagawa M, Ishida S, Suzuki A, Yoshida K, and Japan Levetiracetam N01221 Study Group: Efficacy and tolerability of levetiracetam as adjunctive therapy in Japanese patients with uncontrolled partial-onset seizures. *Psychiatry Clin Neurosci* 2015, 69: 640-8.
18. Fumuro T, Matsumoto R, Shimotake A, Matsuhashi M, Inouchi M, Urayama S, Sawamoto N, Fukuyama H, Takahashi R, Ikeda A: Network hyperexcitability in a patient with partial reading epilepsy: Converging evidence from magnetoencephalography, diffusion tractography, and functional magnetic resonance imaging. *Clin Neurophysiol* 2015, 126: 675-81.
19. Shibata S, Kunieda T, Inano R, Sawada M, Yamao Y, Kikuchi T, Matsumoto R, Ikeda A, Takahashi R, Mikuni N, Takahashi J, Miyamoto S: Risk factors for infective complications with long term subdural electrode implantation in patients with medically intractable partial epilepsy. *World Neurosurg* 2015 84(2): 320-6.
20. Hashi S, Yano I, Shibata M, Masuda S, Kinoshita M, Matsumoto R, Ikeda A, Takahashi R, Matsubara K: Effect of CYP2C19 polymorphisms on the clinical outcome of low-dose clobazam therapy in Japanese patients with epilepsy. *Eur J Clin Pharmacol* 2015, 71: 47-59.
21. Kunieda T, Yamao Y, Kikuchi T, Matsumoto R: New Approach for Exploring Cerebral Functional Connectivity: Review of CCEP (Cortico-cortical evoked potential). *Neurol Med Chir* 2015, 55: 374-382.
22. Matsumoto R, Mikuni N, Tanaka K, Usami K, Fukao K, Kunieda T, Takahashi Y, Miyamoto S, Fukuyama H, Takahashi R, Ikeda A: Possible induction of multiple seizure foci due to parietal tumour and anti-NMDAR antibody. *Epileptic Disord* 2015, 17: 89-94.

〈和文 Japanese articles〉

23. Nakatani M, Tukino M, Takahashi R, Ikeda A : バラシクロビルによる一過性の周期性同期性放電を伴う薬剤性脳症をきたした高齢者例, 臨床神経 2016; 56: 504-507.
24. 井上岳司, 池田昭夫 : 問題症例の脳波ー頭皮上脳波で陽性棘波 (positive spike) を示す症例の病態ー, 臨床神經生理学 2015, 43(6): 477-482.

編集書籍・雑誌 Edited books & Journal

1. 兼本浩祐 主編集, 池田昭夫ら副編集: 臨床てんかん学. 東京. 医学書院, 総ページ数: 671p., 医学書院 2015.

書籍 Book chapters

〈英文 English articles〉

1. Ikeda A: Subdural EEG in frontal lobe epilepsy. Invasive Studies of the Human Epileptic Brain: Principles and Practice of Invasive Brain Recordings and Stimulation in Epilepsy., Oxford University Press, London, 2015, in press.
2. Matsumoto R, Kunieda T: Cortico-cortical evoked potential mapping (Section 5: Human Brain Mapping). In: Lhatoo S, Kahane P, Lüders H, eds. Invasive Studies of the Human Epileptic Brain: Principles and Practice of Invasive Brain Recordings and Stimulation in Epilepsy., Oxford University Press, London, 2015, in press.

〈和文 Japanese articles〉

3. 塚田剛史, 井上岳司, 池田昭夫: てんかん, 神経疾患, 内科診療実践マニュアル第2版, 日本臨床内科医会編, 日本医学出版, 東京, 2016 (印刷中).
4. 藤井大樹, 池田昭夫: 失神, 痙攣, 症候編, 内科診療実践マニュアル第2版, 日本臨床内科医会編, 日本医学出版, 東京, 2016 (印刷中).
5. 吉村元, 池田昭夫: てんかん, JMEDJ 治療法便覧 2016 ~私の治療~, 猿田享男, 北村総一郎総監修, 水澤英洋 神経分野編, 日本医事新報社, 東京, 2016 (印刷中).
6. 池田昭夫, 田中達也: 国際関係, てんかん医療の世界的動向と, 国際組織の中の日本てんかん学会, てんかん白書, 日本てんかん学会てんかん白書編集委員会編, 南江堂 (東京), 2016 (印刷中).
7. 池田昭夫: 専門職 (医師と医学生), てんかん白書, 日本てんかん学会てんかん白書編集委員会編, 南江堂 (東京), 2016 (印刷中).
8. 田中達也, 池田昭夫: 世界のてんかんの潮流, 日本てんかん学会50周年記念誌, 50周年記念誌編集委員会編, 診断と治療社 (東京), 2016 (印刷中).
9. 田中達也, 池田昭夫: 生理的検査と画像の歴史, 日本てんかん学会50周年記念誌, 50周年記念誌編集委員会編, 診断と治療社 (東京), 2016 (印刷中).
10. 村井智彦, 人見健文, 池田昭夫: 進行性ミオクローネスてんかん一成人, 稀少てんかんの診療指標, 井上有史他編, 診断と治療社 (東京), 2016 (印刷中).
11. 坂本光弘, 松本理器, 池田昭夫: 自己免疫介在性脳炎・脳症, 稀少てんかんの診療指標, 井上有史他編, 診断と治療社 (東京), 2016 (印刷中).
12. 下竹昭寛, 池田昭夫: 第1章. 脳波. 6. てんかん. a. 総論, 飛松省三 (編) ここが知りたい! 臨床神経生理, 中外医学社, 東京, 2016, 34-36 (225).
13. 十川純平, 松本理器: 第1章. 脳波. 10. 意識障害, 飛松省三 (編) ここが知りたい! 臨床神経生理, 中外医学社, 東京, 2016, 62-65 (225).
14. 大封昌子, 松本理器: 第1章. 脳波. 11. 脳死, 飛松省三 (編) ここが知りたい! 臨床神経生理, 中外医学社, 東京, 2016, 66-68 (225).

15. 井上岳司, 池田昭夫 : 第1章. 脳波. 14. 脳波レポート (作成・判読所見), 飛松省三(編) ここが知りたい! 臨床神経生理, 中外医学社, 東京, 2016, 75-78 (225).
16. 井上岳司, 池田昭夫 : てんかん外来, II 外来診察の基本的な流れ, 鑑別診断, in:辻貞俊 (Ed.), 神経内科外来シリーズ4, メジカルビュー社, 東京, 2016, 46-52 (232).
17. 人見健文, 寺田清人, 池田昭夫 : 第9章. ミオクローヌス. 第1部. 不随意運動, 不随意運動の診断と治療 改訂第2版, 診断と治療社, 東京, 2016 (印刷中).
18. 吉村元, 松本理器 : II. 外来診療の基本的な流れ. 診断に必要な検査., 辻貞俊(編) 神経内科シリーズ4. てんかん外来, メジカルビュー社, 東京, 2016, 36-45 (232).
19. 小林勝哉, 人見健文, 松本理器, 池田昭夫 : てんかんにおける Borderzone, 中外医学社, 東京, 2016.
20. 村井智彦, 井上岳司, 池田昭夫 : 第7章 てんかん 1 包括的な治療指針, EBMに基づく脳神経疾患の基本治療指針 第4版, メジカルビュー社, 東京, 2016, 370-384 (807).

21. 井上岳司, 人見健文 : 臨床てんかん学 (チャプターナンバー: 1. シアリドーシス. I. 進行性ミオクローヌステンカン. 第11章てんかんおよびてんかん類似症候群). 東京. 医学書院. 403-4, 2015.
22. 井上岳司, 人見健文 : 臨床てんかん学 (チャプターナンバー: 2. ゴーシエ病. I. 進行性ミオクローヌステンカン. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院. 405, 2015.
23. 人見健文, 池田昭夫 : 神経疾患最新の治療2015-2017 (チャプターナンバー: 本態性振戦). 東京. 南江堂. 171-173, 2015.
24. 太田真紀子, 中野智 : 免疫性神経疾患 (チャプターナンバー: その他の免疫性神経疾患 膜原病および類縁疾患に伴う神経筋障害 Susac 症候群). 大阪. 日本臨床社. 716-721, 2015.
25. 石田紗恵子, 池田昭夫 : 臨床てんかん学 (チャプターナンバー: 13. さまざまな焦点を伴う家族性部分てんかん. L. 遺伝子変位がてんかんだけでなく脳症も生ずるか, 遺伝子変異が脳の形態異常をもたらしそれが間接的にてんかんの原因となる疾患. 第11章てんかんおよびてんかん類似症候群). 東京. 医学書院. 451-452, 2015.
26. 下竹昭寛, 松本理器 : 臨床てんかん学 (チャプターナンバー: 5. 頭頂葉てんかん. D. 年齢非依存性焦点性てんかん. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院. 378-380, 2015.
27. 井内盛遠, 松本理器 : 臨床てんかん学 (チャプターナンバー: 1. 側頭葉てんかん. D. 年齢非依存性焦点性てんかん. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院. 367-370, 2015.
28. 人見健文, 池田昭夫 : 臨床てんかん学 (チャプターナンバー: 8. 良性成人型家族性ミオクローヌステンカン. I. 進行性ミオクローヌステンカン. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院. 411, 2015.
29. 十川純平, 人見健文 : 臨床てんかん学 (チャプターナンバー: 7. DRPLA. I. 進行性ミオクローヌステンカン. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院, 410, 2015.
30. 太田真紀子, 人見健文 : 臨床てんかん学 (チャプターナンバー: 4. Lafora 病. I. 進行性ミオクローヌステンカン. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院, 407, 2015.
31. 宇佐美清英, 松本理器 : 臨床てんかん学 (チャプターナンバー: 2. 前頭葉てんかん (ジャクソン発作を除く). D. 年齢非依存性焦点性てんかん. 第11章てんかんおよびてんかん類似症候群). 東京. 医学書院, 371-373, 2015.
32. 小林勝哉, 人見健文 : 臨床てんかん学 (チャプターナンバー: 5. セロイドリポフスチン病. I. 進行性ミオクローヌステンカン. 第11章てんかんおよびてんかん類似症候群.) 東京. 医学書院, 408, 2015.
33. 小林勝哉, 人見健文 : 臨床てんかん学 (チャプターナンバー: 6. Unverricht-Lundborg 病. I. 進行性ミオクローヌステンカン. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院, 409, 2015.

34. 村井智彦, 人見健文: 臨床てんかん学(チャプターナンバー: 3. 赤色ぼろ線維・ミオクローヌスてんかん症候群(MERRF)). I. 進行性ミオクローヌスてんかん. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院, 406, 2015.
35. 芝田純也, 松本理器: 臨床てんかん学 (チャプターナンバー: 4. Jackson 発作関連てんかん. D. 年齢非依存性焦点性てんかん. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院, 376-377, 2015.
36. 金澤恭子, 松本理器: 臨床てんかん学 (チャプターナンバー: 3. 後頭葉てんかん. D. 年齢非依存性焦点性てんかん. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院, 374-375, 2015.
37. 麓直浩, 池田昭夫: 臨床てんかん学 (チャプターナンバー: 12. 家族性側頭葉てんかん. L. 遺伝子変位がてんかんだけでなく脳症も生ずるか, 遺伝子変異が脳の形態異常をもたらしそれが間接的にてんかんの原因となる疾患. 第11章てんかんおよびてんかん類似症候群.). 東京. 医学書院, 451, 2015.
38. 池田昭夫: 臨床てんかん学(チャプターナンバー: B. 局在論からみたてんかん発作. 第7章てんかん発作の症候学.). 東京. 医学書院, 137-141, 2015.
39. 池田昭夫, 赤松直樹, 小林勝弘, 酒田あゆみ, 末永和榮, 飛松省三, 橋本修治, 松浦雅人: 臨床神経生理学 (チャプターナンバー: 日本臨床神経生理学会ペーパレス脳波の記録・判読指針小委員会: デジタル脳波の記録・判読指針). 東京. 臨床神経生理学会, 22-62, 2015, 43.
40. 人見健文, 池田昭夫: デジタル脳波記録・判読の手引き (池田昭夫主編集: 松本理器, 人見健文副編集). (1記録, 第2部指針に基づいた実例提示). 東京. 診断と治療社. 50-54, 2015, Oct.

総説 Review papers

〈英文 English review paper〉

1. Kinoshita M, Ikeda A: Phantom of oscillation: Operational definition bound to improve. editorial. *Clin Neurophysiol*, 2016, 127(1):8-9.
2. Ikeda A: Commentary: We shall not lose past medical documents, *Epilepsia* 2015, 56(11): 1712-3., doi: 10.1111/epi.13196.

〈和文 Japanese review papers〉

3. 音成秀一郎, 池田昭夫: てんかんの診断, 実践! 神経救急 (neurocritical care), 診断と治療』特集 2016 (印刷中).
4. 太田真紀子, 人見健文, 池田昭夫: 特発性全般てんかん, 神経疾患治療ノート, Clinical Neuroscience, 2016 (印刷中).
5. 池田昭夫: てんかんの診断と病型分類, てんかん: 内科医が知っておくべき診療ポイントと治療の最前線, 日本国内科学会雑誌 105 ; 2016 (印刷中).
6. 音成秀一郎, 池田昭夫: 本邦における高齢者てんかんの臨床的特徴, 新薬と臨牀. 65巻 6号 840-845, 2016.
7. 井内盛遠, 池田昭夫: wide-band EEG の時間周波数解析, 目で見るてんかん, Epilepsy 10 : 4-7, 2016.
8. 下竹昭寛, 國枝武治, 松本理器: 皮質脳波記録, Clinical Neurosciece, 2016. 34 (7) : 771-776.
9. 十川純平, 松本理器, 池田昭夫: てんかん病態下の脳内ネットワーク, Clinical Neuroscience, 中外医学社, 東京, 2016, 34 (6) : 713-716.
10. 松本理器, 下竹昭寛, 山尾幸広, 國枝武治: 特集: 神経生理最前線てんかん外科における術前皮質・白質機能マッピング, 最新精神医学, 世論時報社, 東京, 2016, 21 (2) : 101-109.

11. 山尾幸広, 國枝武治, 松本理器: 皮質電気刺激によるヒト脳機能ネットワークの探索, 脳神経外科速報, メディカ出版, 大阪, 2016 Vol. 25, No. 5 in press.
12. 池田昭夫: てんかん発作の発現機構:red slowはあるか?, 脳神経外科ジャーナル, 日本脳神経外科コングレス・三輪書店, 東京, 2016, 128-136.
13. 池田昭夫: てんかん診療を考える, 正しい診断と治療のために, クレデンシャル, 株式会社日本アルトマーク, 東京, 2016, 5-12.
14. 池田昭夫: AES2015印象記, 第69回米国てんかん学会記録集, 2016.
15. 池田昭夫: 編集後記, 臨床神経学, 2016/4/1.
16. 藤井大樹, 池田昭夫: 特集「これからのがん医療～ペランパネルへの期待～」AMPA受容体とがん原性(病態), クリニシアン, エーザイ, 東京, 2016, 63 (5-6) : 29-35.
17. 武山博文, 松本理器, 池田昭夫: くすぶり型辺縁系脳炎と自律神経, 神経内科, 科学評論社, 東京, 2016, Jan, 84 (1) : 58-61.
18. 井上岳司, 池田昭夫: てんかん治療ガイドライン2010, BRAIN NURSING, メディカ出版, 大阪, 2015, 11, 41-45.
19. 池田昭夫: てんかん性DC電位(緩電位):総論「てんかん性wide-band EEG記録と解析のコンセンサスへ:DC電位(緩電位)と高周波振動(HFO)」, 日本臨床神経生理学, 日本臨床神経生理学会, 東京, 2015, Dec, 43 (6) : 482-488.
20. 金澤恭子, 池田昭夫: 発作時DC電位の実際の記録・解析, 「てんかん性wide-band EEGの記録と解析のコンセンサスへ:DC電位(緩電位)と高周波振動(HFO)」日本臨床神経生理学, 日本臨床神経生理学会, 東京, 2015, Dec, 43 (6) : 489-496.

学会・研究会発表 Presentations

学会発表 Congress presentations

■招請講演・シンポジウムなど Invited lectures and symposium etc.

〈国際学会 International presentations〉

1. Ikeda A: Advanced techniques in HFO analysis. (招待講演) Workshop HFOs and the Epileptogenic Zone, 11th Asian & Oceanian Epilepsy Congress (AOEC) (2016/05/13-16, Hongkong, China)
2. Ikeda A: Exploring the mysteries of EEG: Can infraslow and DC shift improve epilepsy treatment? (招待講演) The Masakazu Seino Memorial Lecture, 11th Asian & Oceanian Epilepsy Congress (AOEC) (2016/05/13-16, Hongkong, China)
3. Ikeda A: HFO synchronization and epilepsy network, Parallel session Stereo-EEG and Brain Networks (招待講演) 11th Asian & Oceanian Epilepsy Congress (AOEC) (2016/05/13-16, Hongkong, China)
4. Ikeda A: ictal DC and HFO, and beyond that? Is it red slow? (招待講演) 2nd International Workshop on High Frequency Oscillations in Epilepsy (2016/3/10-12, Freiburg, Germany)
5. Ikeda A: Ictal and interictal infraslow with HFO in human epilepsy (招待講演) Fondazione IECCS Instituto Neurologico Carlo Besta (2016/3/8, Milano, Italy)
6. Hitomi T, Kobayashi K, Takahashi R, Ikeda A: Updates of Neurophysiological Biomarkers in the Genetic Generalized Epilepsy, The 9th KES-JES Joint Symposium, 21st Korean Epilepsy Congress, 2016/6/17-18, Seoul, South Korea.
7. Ikeda A: Wide-band EEG in clinical epilepsy: slow shifts and HFO, How to record and analyze slow shifts in epilepsy (招待講演) 15th European Congress on Clinical Neurophysiology (2015/9/30-10/4, Brno, Czech Republic)
8. Ikeda A: Ictal and interictal slow combined with HFO: Is it red slow? (招待講演) Beijing Epilepsy and Neurophysiology Summit (BENS 2015) (2015/9/12, Beijing, China)
9. Ikeda A: Non-structural evidence of seizure focus: The role of functional imaging, Plenary session Epilepsy Imaging, 31st International Epilepsy Congress, 2015 (2015/9/5-9, Istanbul, Turkey)
10. Matsumoto R: The ventral language network - its function, connectivity and plasticity: insights from direct cortical stimulation and ECoG recordings., Special Seminar at Cognitive Neurophysiology and Brain-Machine Interface Laboratory Department of Neurology, Johns Hopkins Hospital (2015/12/8, Baltimore, USA)

〈国内学会 Domestic presentations〉

11. 池田昭夫：てんかん原性の臨床生理学的 biomarker : red slow はあるか？, 第12回日本てんかん学会近畿地方会, 大阪, 2016/7/23.
12. 池田昭夫：Hands-on 3 「脳波」, 日本神経学会学術大会 第13回生涯教育セミナー, 神戸, 2016/5/21.
13. 松本理器：Hands-on 3 「脳波」, 日本神経学会学術大会 第13回生涯教育セミナー, 神戸, 2016/5/21.
14. 池田昭夫：Uncommon, but important causes of epilepsy and seizure, 第57回日本神経学会学術大会 International Educational Session 03 How to Manage Seizures, 神戸, 2016/5/18-21.
15. 松本理器：自己免疫性てんかんとの関連, 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.

16. 松本理器：非けいれん性てんかん重積(NSCE)：診断と治療，第57回日本神経学会学術大会，神戸，2016/5/18-21.
17. 池田昭夫：オシロロジーからみた神経疾患の病態と治療，第57回日本神経学会学術大会（シンポジウム 成人てんかんとオシロロジー），神戸，2016/5/18-21.
18. 吉村元，松本理器，池田昭夫，幸原伸夫：高齢者てんかん重積状態の臨床的特徴と治療（シンポジウム 治るてんかん：高齢者てんかんの診断と治療），第57回日本神経学会学術大会，神戸，2016/5/18-21.

19. 池田昭夫：成人てんかん治療：薬剤と生理学手法の可能性（招待講演）第42回日本脳科学学会てんかんシンポジウム宮崎2015，宮崎，2015/11/13.
20. 小林勝哉，松本理器，國枝武治，池田昭夫：慢性電極留置におけるてんかん原性評価，第45回日本臨床神経生理学会学術大会，大阪，2015/11/5-7.
21. 池田昭夫：臨床神経生理学的所見とグリア機能のオーバービュー（シンポジウム3），第49回日本てんかん学会学術集会，長崎，2015/10/30-31.
22. 池田昭夫，Bayasgalan Borgil，松橋眞生，文室知之，中野直樹，飯田幸治，片桐匡弥，松本理器，國枝武治，加藤天美，高橋良輔：VNS の効果作動機構の検討（シンポジウム4），第49回日本てんかん学会学術集会，長崎，2015/10/30-31.
23. 小林勝哉，松本理器，宇佐美清英，國枝武治，池田昭夫：睡眠と前頭葉てんかん，第49回日本てんかん学会学術大会，長崎，2015/10/30-31.

■一般発表 Oral and poster presentations

国際学会 International presentations

1. Daifu M, Inouchi M, Kanazawa K, Matsuhashi M, Inoue T, Kobayashi K, Shimotake A, Hitomi T, Matsumoto R, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Intracranially recorded ictal DC shifts and HFOs as surrogate markers of epileptogenicity in human focal epilepsy, 2nd International Workshop on High Frequency Oscillations in Epilepsy. (2016/3/10-12 Freiburg, Germany)

2. Yoshimura H, Matsumoto R, Ishii J, Kono T, Hoshi T, Todo K, Kawamoto M, Ariyoshi K, Ikeda A, Kohara N: Status epilepticus in the elderly in the super-aging society: clinical and EEG features and prognostic factors, 31st International Epilepsy Congress, 2015 (2015/9/5-9, Istanbul, Turkey)

国内学会 Domestic presentations

〈英語発表 English presentations〉

3. Daifu M, Matsumoto R, Shimotake A, Sakamoto M, Ota M, Kunieda T, Miyamoto S, Takahashi R, Ralph MAL, Ikeda A: Ventral part of the anterior temporal lobe actively engages in Kanji reading: Evidence from direct cortical stimulation of the basal temporal language area. 第39回日本神経科学大会，横浜，パシフィコ横浜，2016/7/20-22.
4. Shimotake A, Matsumoto R, Kobayashi K, Kunieda T, Mikuni N, Miyamoto S, Takahashi R, Ikeda A: 2016 7/20-22. Functional mapping of praxis: Electrical cortical stimulation study. 第39回神経科学大会. パシフィコ横浜., 2016/7/20-22.

5. Takeyama H, Matsumoto R, Usami K, Shimotake A, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Human motor inhibition network as revealed by epicortical ERP, CCEP and high frequency cortical stimulation. 第39回日本神経科学大会, 横浜, パシフィコ横浜, 2016/7/22.
6. Nakae T, Matsumoto R, Kunieda T, Arakawa Y, Kobayashi T, Inada T, Takahashi Y, Nishida S, Inano R, Kobayashi K, Shimotake A, Matsuhashi M, Kikuchi T, Takahashi R, Ikeda A, Miyamoto S: Antero-posterior gradients exist in functional connections between human inferior frontal gyrus and middle&posterior temporal gyri. An intraoperative cortico-cortical evoked potential study. 第39回日本神経科学大会, 横浜, 2016/7/20-22.
7. Togo M, Matsumoto R, Takeyama H, Kobayashi K, Shimotake A, Usami K, Inouchi M, Nakae T, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Functional connectivity from medial parietal cortex and superior parietal lobule: a cortico-cortical evoked potential study. 第39回日本神経科学大会, 横浜, 2016/7/20-22.
8. Ota M, Shimotake A, Matsumoto R, Sakamoto M, Daifu M, Nakae T, Kunieda T, Miyamoto S, Takahashi R, M.Lambon-Ralph, Ikeda A: Compensatory mechanisms of semantic- and kanji / kana word processing after resection of the anterior temporal lobe (in epilepsy surgery). 第39回日本神経科学大会, 横浜, 2016/07/20-22.
9. Kinboshi M, Mukai T, Tokudome K, Kunisawa N, Higor A. Iha, Shimizu S, Serikawa T, Ito H, Ikeda A, Ohno Y: Expressional analysis of astrocytic Kir4.1 channels in audiogenic seizure-susceptible Lgi1L385R/+ mutant rats. 第39回日本神経科学大会, 横浜, 2016/07/20-22.
10. Yoshinaga K, Matsuhashi M, Hanakawa T, Ikeda A: Phase angle has information on temporal relationship of signals in functional connectivity study in human brain magnetoencephalography analysis. 第39回日本神経科学大会, 横浜, 2016/07/20-22.
11. Nakatani M, Matsumoto R, Kobayashi K, Hitomi T, Inouchi M, Matsuhashi M, Kinoshita M, Kunieda T, Miyamoto S, Hattori N, Takahashi R, Ikeda A: Repeated, cortical stimulation inhibits epileptic excitability in human epileptic focus. 第57回日本神経学会学術大会, 神戸, 2016/5/18-5/21.
12. Takeyama H, Matsumoto M, Usami K, Shimotake A, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Which areas inhibit motor responses? Direct evidence from subdural recordings. 第57回日本神経学会学術大会, 神戸, 神戸国際会議場, 2016/5/18-21.
13. Togawa J, Inouchi M, Matsumoto R, Matsuhashi M, Kobayashi K, Usami K, Hitomi T, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Enhancement of coupling between slow waves and high gamma activities during slow wave sleep in human. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
14. Daifu M, Inouchi M, Kanazawa K, Matsuhashi M, Inoue T, Kobayashi K, Shimotake A, Hitomi T, Matsumoto R, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Surrogate marker of intracranially recorded ictal DC shifts and HFOs in human focal epilepsy. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
15. Neshige S, Sakurai T, Kobayashi K, Shimotake A, Matsuhashi M, Matsumoto R, Hitomi T, Matsumoto M, Takahashi R, Ikeda A: Wide-band Electrocorticographic (ECOG) data analysis in primary sensorimotor (SI-MI) area. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
16. 松本理器：自己免疫性てんかんとの関連, 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
17. Shimotake A, Matsumoto R, Togo M, Kunieda T, Ikeda A: Functional mapping of praxis network: Electrical cortical stimulation study. (ポスター) The 1st International Symposium on Embodied-Brain Systems Science (EMBOSS2016), 東京, 2016/5/8-9.
18. Sakamoto M, Matsumoto R, Shimotake A, Nakae T, Kunieda T, Ikeda A: Role of left ventral precentral gyrus in apraxia of speech: documents with electrical cortical stimulation and focal resection. (ポスター) The 1st International Symposium on Embodied-Brain Systems Science(EMBOSS2016), 東京, 2016/5/8-9.

19. Togo M, Matsumoto R, Shimotake A, Inouchi M, Kunieda T, Ikeda A: Functional connectivity from human superior parietal lobule and medial parietal lobe: a cortico-cortical evoked potential (CCEP) study. (ポスター) The 1st International Symposium on Embodied-Brain Systems Science (EMBOSS2016), 東京, 2016/5/8-9.
20. Daifu M, Inouchi M, Inoue T, Kanazawa K, Matsuhashi M, Kobayashi K, Shimotake A, Hitomi T, Matsumoto R, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Co-occurrence of slow shifts and high frequency oscillations (HFOs) in invasively recorded, interictal state: Is it called 'Red slow'? 第49回日本てんかん学会学術集会, 長崎, 2015/10/30-31.
21. Nakatani M, Matsumoto R, Kobayashi K, Hitomi T, Inouchi M, Matsuhashi M, Kinoshita M, Kunieda T, Miyamoto S, Hattori N, Takahashi R, Ikeda A: Modulation of wide-band electrocorticographic activities by electric cortical stimulation. 第49回日本てんかん学会学術集会, 長崎, 2015/10/30-31.
22. Murai T, Hitomi T, Inoue T, Kobayashi K, Shimotake A, Matsuhashi M, Inouchi M, Matsumoto R, Takahashi R, Ikeda A: Analysis of scalp-recorded, ictal direct current shift: its sensitivity and specificity. 第49回日本てんかん学会学術集会, 長崎, 2015/10/30-31.
23. Takeyama H, Matsumoto R, Kobayashi K, Usami K, Shimotake A, Kikuchi T, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Functional connectivity of the human entorhinal cortex: a cortico-cortical evoked potential study. 第49回日本てんかん学会学術集会, 長崎, 2015/10/30-31.

〈日本語発表 Japanese presentations〉

24. 本多正幸, 下竹昭寛, 坂戸勇介, 引網亮太, 藤井大樹, 小林勝哉, 松本理器, 池田昭夫, 高橋良輔: 食事中の大きな笑いにより強直発作様の意識消失を呈した73歳男性例. 第12回 日本てんかん学会近畿地方会, 大阪, 2016/7/23.
25. 坂本光弘, 松本理器, 十川純平, 武山博文, 端祐一郎, 小林勝哉, 下竹昭寛, 近藤誉之, 高橋良輔, 池田昭夫: 自己免疫機序が関与するてんかん症候群の診断アルゴリズム構築の試み. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
26. 村井智彦, 人見健文, 小林勝哉, 下竹昭寛, 岡田知久, 松本理器, 富樫かおり, 澤本伸克, 高橋良輔, 池田昭夫: BAFMEにおけるPET画像上の特徴. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
27. 谷岡洸介, 人見健文, 松本理器, 飛松省三, 犬塚貴, 吉良潤一, 楠進, 高橋良輔, 池田昭夫: 脳波判読セミナーのアンケート調査からの脳波教育の現状解析;過去5年間の実態とニーズ. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
28. 人見健文, 小林勝哉, 井内盛遠, 櫻井健世, Shamima Sultana, 佐藤啓, 井上岳司, 下竹昭寛, 松本理器, 陳和夫, 高橋良輔, 池田昭夫: 良性成人型家族性ミオクローネスてんかんにおけるてんかん性放電の睡眠による変容. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
29. 十河正弥, 人見健文, 村井智彦, 松橋真生, 松本理器, 幸原伸夫, 高橋良輔, 池田昭夫: 低酸素脳症の頭皮上脳波における脳波所見と超低周波活動 (infraslow activity) の検討. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
30. 端祐一郎, 坂本光弘, 十川純平, 岡田洋一郎, 藤井ちひろ, 越智博文, 中川正法, 水野敏樹, 下竹昭寛, 松本理器, 漆谷真, 池田昭夫, 高橋良輔, 近藤誉之: 自己免疫性てんかんにおける末梢血リンパ球の表面抗原解析. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.
31. 三橋賢大, 北村彰浩, 小林勝哉, 上村紀仁, 葛谷聰, 山下博文, 澤本伸克, 近藤誉之, 漆谷真, 池田昭夫, 高橋良輔: 脊髄サルコイドーシスの治療に関する臨床的検討. 第57回日本神経学会学術大会, 神戸, 2016/5/18-21.

32. 岡田信久, 和田隆史, 樽野陽亮, 真木崇州, 吉藤元, 丹治正大, 三橋賢大, 山門穂高, 上田紗希帆, 前川嵩太, 引網亮太, 木下久徳, 綾木孝, 山下博文, 澤本伸克, 漆谷真, 池田昭夫, 高橋良輔: 再発性多発軟骨炎に伴う脳炎2症例の検討, 第57回日本神経学会学術大会. 神戸国際会議場. 2016/5/18-21.
33. 佐藤啓, 人見健文, 葛谷聰, 武地一, 木下彩栄, 池田昭夫: 高橋良輔: 認知症鑑別診断への脳波利用の試み, 第113回日本内科学会総会, 東京, 2016/4/15-17.
34. 松本理器, 國枝武治, 池田昭夫: 皮質・皮質間誘発電位 (CCEP) 記録・解析の実際, ワークショップ4. 皮質・皮質間誘発電位, 第45回日本臨床神経生理学会学術大会, 大阪, 2015/11/5-7.
35. 人見健文, 小林勝哉, 櫻井健世, Shamima Sultana, 佐藤啓, 井上岳司, 下竹明寛, 松本理器, 高橋良輔, 池田昭夫: 睡眠が良性成人型家族性ミオクローヌスてんかん (BAFME) のてんかん性放電に与える影響, 第45回日本臨床神経生理学会学術大会, 大阪, 2015/11/5-7.
36. 十河正弥, 松本理器, 武山博文, 小林勝哉, 下竹昭寛, 宇佐美清英, 中江卓郎, 國枝武治, 宮本享, 高橋良輔, 池田昭夫: ヒト内側頭頂葉の機能的結合性: 皮質-皮質間誘発電位を用いた検討, 第45回日本臨床神経生理学会学術大会, 大阪, 2015/11/5-7.
37. 中江卓郎, 松本理器, 國枝武治, 荒川芳輝, 下竹昭寛, 小林勝哉, 稲野理賀, 西田誠, 高橋由紀, 稲田拓, 菊池隆幸, 松橋眞生, 高橋良輔, 池田昭夫, 宮本享: 術中電気的線維追跡法を用いた腹側言語経路の検討, 第45回日本臨床神経生理学会学術大会, 大阪 (大阪国際会議場), 2015/11/5-7.
38. 下竹昭寛, 松本理器, 太田真紀子, 國枝武治, 宮本享, 高橋良輔, 池田昭夫: 側頭葉底部前方領域での意味認知機能マッピング: 事象関連電位の皮質記録の検討, 第49回日本てんかん学会学術集会, 長崎, 2015/10/30-31.
39. 人見健文, 小林勝哉, 高橋良輔, 池田昭夫: 良性成人型家族性ミオクローヌスてんかん (BAFME) の発症の予測因子の検討, 第49回日本てんかん学会学術集会, 長崎, 2015/10/30-31.
40. 井上岳司, 酒井達也, 小林勝哉, 下竹昭寛, 井内盛遠, 松本理器, 池田昭夫, 高橋良輔: Levetiracetam により paradoxical effect を認めた3例の臨床的検討, 第49回日本てんかん学会総会, 長崎, 2015/10/30-31.
41. 吉村元, 松本理器, 上田浩之, 藤原悟, 上田哲大, 引網亮太, 村瀬翔, 石井淳子, 河野智之, 星拓, 藤堂謙一, 川本未知, 有吉孝一, 池田昭夫, 幸原伸夫: 高齢者のてんかん重積状態における頭部MRI拡散強調画像: 臨床・脳波所見との相関, 第49回日本てんかん学会学術集会, 長崎, 2015/10/30-31.
42. 中江卓郎, 國枝武治, 松本理器, 荒川芳輝, 西田誠, 稲野理賀, 高橋由紀, 稲田拓, 下竹昭寛, 菊池隆幸, 吉田和道, 高木康志, 池田昭夫, 宮本享: 意味認知の機能可塑性: 側頭葉腫瘍切除症例における神経心理学的検討, 日本脳神経外科学会第74回学術総会, 札幌 (さっぽろ芸術文化の館), 2015/10/14-16.
43. 谷岡洸介, 丸本圭一, 魚住容子, 谷口浩一郎, 濱野利明, 立花直子: 閉塞性無呼吸症候群患者における口腔内装置治療後の自覚症状改善因子, 第7回日本臨床睡眠医学会, 大阪, 2015/8/1.

その他研究会など Other presentations

〈英語発表 English presentations〉

1. Togo M, Matsumoto R, Shimotake A, Inouchi M, Kunieda T, Ikeda A: Functional connectivity from human superior parietal lobule and medial parietal lobe: a cortico-cortical evoked potential (CCEP) study. 第3回身体性システム領域全体会議, 花巻, 2016/3/7-9.
2. Togawa J, Inouchi M, Matsumoto R, Matsuhashi M, Kobayashi K, Usami K, Hitomi T, Kunieda T, Miyamoto S, Takahashi R, Ikeda A: Dynamic alteration of coupling between slow waves and high gamma activities in different sleep stages: human electrocorticogram (ECoG) study. 神経データ解析ワークショップ, 京都, 2016/2/22.

〈日本語発表 Japanese presentations〉

3. 池田昭夫：側頭葉由来の新しいてんかん症候群（扁桃体腫大，免疫性くすぶり型辺縁系脳炎，家族性外側側頭葉てんかんなど）（講演），第13回広島けいれんフォーラム，広島，2016/7/7.
4. 池田昭夫：てんかん発作とてんかんの治療 急性期から慢性期まで（講演），第5回 Kanto Epilepsy Plaza，東京，2016/5/27.
5. 池田昭夫：てんかん原性の臨床生理学的 biomarker ? – Red slow はあるかー（講演），東海神経生理懇話会，名古屋，2016/4/23.
6. 池田昭夫：てんかん診療の実際 –外来での対応を中心にー（講演），平成27年度後期鹿精協院長等研修会 精神保険指定医研修会，鹿児島，2016/2/25.
7. 池田昭夫：てんかんの診療と治療の基本 新規抗てんかん薬単剤療法のミッション（講演），第3回てんかんの診療連携を考える会，高知，2016/1/22.
8. 池田昭夫：A03：ヒト脳発振現象の直接記録 ヒト脳からの実記録データとその解釈，オシロロジー第1回領域会議，福岡，2016/1/14-15.
9. 松本理器：てんかんの診断と治療の基本（講演），てんかん治療学術講演会，豊田，2016/6/29.
10. 松本理器：Updating in Autoimmune Epilepsy（講演），Novartis Neuro Forum，東京，2016/5/7.
11. 松本理器：てんかんと自己免疫の関連，第9回成人の為のてんかん診療フォーラム，名古屋，2016/4/16.
12. 人見健文：ミオクローヌス（シンポジウム），第31回日本大脳基底核研究会，秋田，2016/7/23-24.
13. 坂本光弘：自己免疫性脳炎と認知症（講演），第2回アルツハイマー病とてんかんを考える会，京都，2016/3/14.
14. 下竹昭寛：挙児希望女性に対するてんかん管理（講演），第6回京滋てんかん病診連携研究会，京都，2016/2/20.
15. 坂本光弘，松本理器，下竹昭寛，國枝武治，池田昭夫：左中心前回中・下部に限局した腫瘍切除後に anarthria を呈した1例（講演），第55回近畿高次神経機能研究会，大阪，2016/2/13.
16. 大封昌子，井内盛遠，松橋眞生，池田昭夫：神経細胞とグリア細胞のてんかん発作発現における機序（講演／口演／ポスター），オシロロジー第1回領域会議，福岡，2016/1/14-15.
17. 池田昭夫：側頭葉由来の新しいてんかん症候群 家族性外側側頭葉てんかん免疫性くすぶり型辺縁系脳炎を中心（講演），第68回大阪小児てんかん研究会，大阪，2015/11/28.
18. 池田昭夫：てんかん原性の臨床生理学的 biomarker ?（講演），第49回埼玉てんかん懇話会，埼玉，2015/11/26.
19. 池田昭夫：側頭葉由来の新しいてんかん症候群（講演），第41回東京てんかんフォーラム，東京，2015/11/20.
20. 松本理器：ヒト脳の解剖・機能的結合地図：非侵襲・侵襲的計測による探索，第27回臨床神経生理研究会，宮崎，2015/8/1-2.

講義, セミナー, 勉強会など Educational lectures

1. 池田昭夫：てんかん性異常と関連脳波所見（講演），第9回脳波・筋電図セミナー，京都，2016/1/30.
2. 池田昭夫：脳波判読（講演），第9回脳波・筋電図セミナー，京都，2016/1/30.
3. 松本理器：非てんかん性異常（講演），第9回脳波・筋電図セミナー，京都，2016/1/30.
4. 池田昭夫：小児難病の視点からのてんかんの診断と治療；最近の話題（講演），第一回難病専門家育成セミナー（徳島大学医学部神経内科），徳島，2015/11/11.
5. 池田昭夫：100人に一人のてんかん 高齢者，認知症，運転免許等をめぐる話題 てんかんの最近の診断と治療（講演），第16回京都府医師会生涯教育セミナー，京都，2015/9/26.
6. 池田昭夫：意識障害・脳波レポートの書き方（講演），第1回脳波セミナー・アドバンスコース，京都，2015/8/9.
7. 松本理器：高齢者のてんかん（講演），第16回京都府医師会生涯教育セミナー，京都，2015/9/26.

学部講義, 院内講義 Intramural lectures

1. 池田昭夫：てんかん総論 ベッドサイド診断（問診のポイントなど）=感度と特異度を常に意識する，脳波・てんかん講義シリーズ，2016/7/28.
2. 池田昭夫：脳のダイナミズムとてんかん発作，神経内科は面白い！2016，2016/7/3.
3. 池田昭夫：脳死移植 脳死判定の実際，第9回チーム移植医療検討会，2016/6/20.
4. 池田昭夫：てんかん・運動異常症の診断と治療，S7ab 臨床神経学（神経内科学・脳神経外科学），2016/5/25.
5. 松本理器：脱髓疾患・自己免疫疾患，S7ab 臨床神経学（神経内科学・脳神経外科学），2016/5/24.
6. 下竹昭寛：脊髄疾患の診断と内科的治療，S7ab 臨床神経学（神経内科学・脳神経外科学），2016/5/26.
7. 人見健文：末梢神経疾患，神経伝導検査と筋電図，S7ab 臨床神経学（神経内科学・脳神経外科学），2016/5/23.
8. 小林勝哉：第13回「てんかんの治療：概論」，神経内科研修セミナー，2016/3/24（木）.

9. 池田昭夫：てんかん診断と治療の基礎，外来ナース講義，2015/8/28.
10. 人見健文：第4回 「脳波の基礎，正常脳波」，脳波・てんかんレクチャー前期，2015/8/20.
11. 人見健文：第4回 「脳波の基礎，正常脳波」，脳波・てんかんレクチャー後期，2015/12/7.
12. 人見健文：脳波：正常所見，人間健康科学系専攻 臨床生理超音波診断学講義，2015/12/25.
13. 小林勝哉：第6回 「てんかん発作重積治療，発作時対応」，脳波・てんかんレクチャー前期，2015/9/3.
14. 小林勝哉：第4回 「てんかん発作重積治療，発作時対応」，脳波・てんかんレクチャー後期，2015/12/10.

VI

Attached materials


Collaborative research activity in the Grant-in-Aid for Scientific Research on Innovative Areas: Non-linear Neuro-oscillology - Towards Integrative Understanding of Human Nature.

This five-year innovative research program is led by Prof. Atsushi Nambu (National Institute for Physiological Sciences, Division of System Neurophysiology) and aims to create a new academic field of Neuro-oscillology, which enables us to understand human nature. We will devote ourselves to researches on wideband-EEG oscillations to understand the human nature and network diseases such as epilepsy (Research Project A03: Principal Investigator Akio Ikeda) and to collaboration with the researchers of other fields including mathematical modeling, intervention and Exploration groups.

Homepage: <http://www.nips.ac.jp/oscillology/>

A03 ヒト脳発振現象の直接記録

ヒト脳機能は多次元・多階層の発振現象の非線形の相互作用により発現する。一方、脳機能の異常発現も作動原理の根本は共通し、てんかんは、自律的な脳ネットワークが突発的に種々の次元・階層で過剰発振する「ネットワーク病」と捉えられる。本研究計画では、正常脳機能およびてんかん発作発現にかかわる局所および広域の集団発振現象をヒト脳からの直接記録で探索する。



空間的観点からは、局所神経回路(細胞外多電極記録)からシステムレベル(皮質脳波、頭皮上脳波、脳磁図)で、発達過程の観点からは、ヒトの乳児・小児・成人脳および動物モデルで比較検討する。B班と連携し、記録データからの数理モデルを構築し、正常振動現象および病態下の異常発振の作動原理と制御機構の解明を目指す。C班と連携し、外的および内的な振動制御の手法を用いて、突発性振動異常の制御を試み、介入による多次元・多階層での生理的・病的振動の変容機構を明らかにする。



研究代表者	池田 昭夫 京都大学大学院 医学研究科 てんかん運動異常生理学
研究分担者	小林 勝弘 岡山大学 医学部 小児神経学
	長峯 隆 札幌医科大学 医学部 神経科学
	松本 理器 京都大学大学院 医学研究科 てんかん運動異常生理学
	國枝 武治 京都大学大学院 医学研究科 脳神経外科学
研究協力者	小林勝哉 京都大学大学院 医学研究科 臨床神経学



2nd group meeting (June 27th-28th, 2016, Sapporo)

9th electroencephalography (EEG) and electromyography (EMG) seminar

January 30th, 2016, Kyoto University

The 1st EEG and EMG seminar was held in February 2008 in Kyoto in order to acquire and improve the basic knowledge and techniques of clinical neurophysiological studies. From the 2nd EEG and EMG seminar, this seminar became a training session related to Japanese Society of Clinical Neurophysiology. Since then, the seminar has been held every winter in Kyoto. Our department served as the secretary office of this seminar. About 150 participants attended comprehensive lectures and hands-on about clinical neurophysiological studies.



1st advance course of electroencephalography (EEG) seminar

August 8th-9th, 2015, Shirankaikan and Shirankaikan Annex

The advance course of electroencephalography (EEG) seminar was founded by Japanese Society of Clinical Neurophysiology (A committee of advance course of electroencephalography (EEG) seminar: Chair Akio Ikeda) in order to acquire specialized knowledge and technique about clinical EEG. The seminar adopted ANZAN (Australian-New Zealand Association of Neurology) style which consists of 8 sessions, and each session consists of lecture (30 min) followed by practical hands-on related to lecture (60 min). About 50 participants deepened their knowledge about recording and reading of clinical EEG.

EEG samples were prepared by the courtesy of ANZAN (Prof. Andrew Bleasel, Prof. Ernest Somerville, Prof. John W. Dunne, and Prof. Nicholas Lawn).



104th Kinki Regional Division of Japanese Society of Neurology

March 6th, 2016, Senri Life Science Center, President Akio Ikeda



Epilepsy Lectures in Autumn 2015, Kyoto

October 26th, 2015, Shirankai Annex

We invited Prof. Solomon Moshé (Albert Einstein College of Medicine, ILAE past president) and Prof. Gary Mathern (UCLA, Epilepsia Chief Editor) to give special lectures before the annual meeting of the Japan Epilepsy Society.



5th Masakazu Seino Memorial Lecture

May 13th, 2016, Hong Kong

Prof. Akio Ikeda gave Masakazu Seino Memorial Lecture in the 11th Asian & Oceanian Epilepsy Congress (AOEC).

5th Masakazu Seino Memorial Lecture

Lecturer:
Prof Akio Ikeda
Kyoto, Japan

**"Exploring the mysteries of EEG:
can infraslow and DC shift
improve epilepsy treatment?"**

Chairman:
Dr Shih-Hui Lim
Singapore

Hong Kong Convention and Exhibition Centre

Visiting physicians

[Overseas]

October 26th, 2015

Prof. Gary Mathern (Epilepsia, Chief editor)

Affiliation: Neurological Surgery, UCLA-David Geffen School of Medicine

Position: Professor

Prof. Solomon Moshé (ILAE, past president)

Affiliation: Charles Frost Chair In Neurosurgery and Neurology

Professor of Neurology, Neuroscience & Pediatrics

Vice-Chair, Dept. of Neurology

Director Pediatric Neurology

Director Clinical Neurophysiology

Albert Einstein College of Medicine and Montefiore Medical Center

January 20th, 2016

Prof. Matthew A. Lambon-Ralph, FRSR (hons), FBPsS

Affiliation: The University of Manchester

Position: Professor of Cognitive Neuroscience & Associate Vice-President (Research)

February 22nd, 2016

Dr. Junji Ito

Affiliation: Institute of Neuroscience and Medicine, Computational and Systems Neuroscience & Institute for Advanced Simulation, Theoretical Neuroscience, Research Centre Jülich, Germany

Position: PostDoc

Prof. Sonja Gruen

Affiliation: Institute of Neuroscience and Medicine, Computational and Systems Neuroscience & Institute for Advanced Simulation, Theoretical Neuroscience, Research Centre Jülich, Germany

Position: Professor

July 20th, 2016

Prof. Michel Thiebaut de Schotten, PhD, CR2 CNRS

Affiliation: Brain Connectivity and Behaviour Group

FrontLab, Brain and Spine Institute

ICM, Hôpital Pitié-Salpêtrière, Paris, FRANCE

Position: Associate Professor

[Domestic]

December 21st, 2015

Yoshio Sakurai, Ph.D

Affiliation: Laboratory of Neural Information, Graduate School of Brain Science, Doshisha University

Position: Professor

February 3rd, 2016

Hiroshi Imamizu, M.D., Ph.D.

Affiliation: Department of Psychology, The University of Tokyo

Position: Professor

VI. Attached materials Visiting physicians

March 15th–16th, 2016

Prof. Ichiro Tsuda, Ph.D.

Affiliation: Department of Mathematics, Hokkaido University

Position: Professor

Prof. Toshio Aoyagi, Ph.D.

Affiliation: Department of Applied Analysis and Complex Dynamical Systems, Kyoto University

Position: Professor

Dr. Keiichi Kitajo, Ph.D.

Affiliation: Rhythm-based Brain Information Processing Unit, BTCC

Position: Unit Leader

Prof. Katsunori Kitano, Ph.D.

Affiliation: Department of Human and Computer Intelligence, Ritsumeikan University

Position: Professor

Yutaka Yamaguti, Ph.D.

Affiliation: Fukuoka Institute of Technology

Position: Assistant Professor

Takao Namiki, Ph.D.

Affiliation: Department of Mathematics, Hokkaido University

Position: Associate Professor

Jiro Okuda, Ph.D.

Affiliation: Department of Intelligent Systems, Faculty of Computer Science and Engineering, Kyoto Sangyo University

Position: Professor

Akihiro Yamaguchi, Ph.D.

Affiliation: Fukuoka Institute of Technology

Position: Associate Professor

Satoru Tadokoro

Affiliation: Department of Mathematics, Hokkaido University

Position: Research Fellow

Keiko Usui, M.D., Ph.D.

Affiliation: Department of Systems Neuroscience, Sapporo Medical University

Position: Lecturer

March 30th, 2016

Kouichi Nakamura, Ph.D.

Affiliation: Department of Morphological Brain Science, Graduate School of Medicine, Kyoto University

Position: Assistant Professor

March 31st, 2016

Kohei Nakajima, Ph.D.

Affiliation: The Hakubi Center for Advanced Research & Graduate School of Informatics, Kyoto University

Position: Assistant Professor

April 6th, 2016

Eiichi Naito, Ph.D.

Affiliation: Brain Networks and Communication Laboratory, Center for Information and Neural Networks (CiNet), National Institute of Information and Communications Technology (NICT)

Position: Research Manager

京都大学大学院医学研究科てんかん・運動異常生理学講座
Department of Epilepsy, Movement Disorders and Physiology
Kyoto University Graduate School of Medicine

年次報告書 Annual Report

発行 2016年8月

発行元 京都大学大学院医学研究科てんかん・運動異常生理学講座
〒606-8507 京都市左京区聖護院川原町 54
TEL : 075-751-3662 FAX : 075-751-3663

印刷 ユニバース印刷
〒620-1441 京都府福知山市三和町梅原 867-1
TEL : 0773-58-2029 FAX : 0773-58-2028
